




**ERNST
SEEDS**



60
YEARS

2024-25 CATALOG



OUR mission

It is the mission of Ernst Seeds to make available to Eastern North America the key native and naturalized species of plants for restoration, reclamation, conservation, wildlife and pollinator habitat enhancement, renewable biomass energy, and beautification of our nation.



Ernst Seeds
8884 Mercer Pike
Meadville, PA 16335

DIRECTIONS

Our farm is located in Northwest Pennsylvania, 80 miles north of Pittsburgh and 40 miles south of Lake Erie off I-79. To get here:

- Take I-79 to Meadville Exit 147A (US 322 East to Meadville).
- Travel on US 322 East to 2nd traffic light.
- Turn right onto Mercer Pike and travel approximately 1,000 feet before turning left to continue on Mercer Pike (follow the sign to the "Ernst Bike Trail").
- Drive two miles on Mercer Pike.
- After crossing the railroad tracks, turn left into the Ernst Lane.

(Note: Some internet and GPS directions are not accurate.)

CONTENTS

Ordering Information	3
A Letter From The President	4
Our Roots Run Deep	6
Meet The Team	8
Index	132
Acknowledgments	136

planting resources

Defining Native Vs. Naturalized	10
Understanding Pure Live Seed (PLS)	11
Expectations Of Native Species	12
Establishment Guide	16
Obstacles To A Successful Meadow Establishment	58
Biomass Overview	64
Bioengineering Materials	72

planting guides

Upland & Meadow Sites	23
Wildlife Habitat & Food Plot Sites	27
Pollinator-Friendly Sites	30
Riparian Sites	33
Disturbed Sites & Steep Slopes	36
Utility Right-Of-Way Sites	40
Wet Meadow & Wetland Sites	42
Partially Shaded Sites	46
Stormwater Management Sites	50
Southeastern U.S. Sites	54
Biomass Production Sites	68

species we offer

Grasses & Grass-Like Species	78
Herbaceous Flowering Species	94
Woody Trees, Shrubs & Vines	119
Cover Crops	125



TAKE THE GUESSWORK OUT OF YOUR PLAN

Knowing what to plant—where, when, and why—is the root of our business. Whether you need help brainstorming options or a pro to stamp your master plan, our team of experts are standing by.

Reach out today, and let's get growing!

FOLLOW US



ORDERING INFORMATION

WAYS TO PLACE YOUR ORDER

Our office hours are Monday through Friday, 8 a.m. to 5 p.m. EST.



TOLL-FREE
(800) 873-3321

PHONE
(814) 336-2404



E-MAIL
sales@ernstseed.com



FAX
(814) 336-5191



ADDRESS
8884 Mercer Pike
Meadville, PA
16335

TERMS & CONDITIONS

ORDER MINIMUMS

- ▶ \$25.00 order minimum before freight and tax.
- ▶ Mixes must be ordered in whole pound increments.
- ▶ 5% custom mix charge.

PAYMENT

Visa, MasterCard, Discover, and American Express are accepted forms of payment. Checks and money orders are also acceptable with placement of order. Sorry, Ernst cannot process COD orders.

FREIGHT

Orders are shipped via UPS or commercial freight (as applicable) and require a street address. Freight is prepaid and added to your invoice unless prior arrangements are made with an Ernst sales associate.

OTHER FEES

Other fees may include, but are not limited to, small order fees, pallet packaging, custom orders, and bioengineering surcharges. We may require payment in advance for certain custom or special orders.

CANCELLATIONS & RETURNS

We cannot guarantee that cancellation requests can be processed within 48 hours of the shipping date. Cancellation requests are subject to Ernst approval.

All plants are guaranteed to be what is specified on the plant tag or shipping documents and healthy upon receipt. Since we cannot guarantee care of such items after they leave our dock, all live plant material claims must be made within 48 hours of receipt.

All orders are subject to acts of natural destruction crop failures; liability is limited to refund or replacement.

Returns must be approved by Ernst and are subject to a restocking fee. Individual items and Ernst stock mixes are subject to a 10% restocking fee and must be returned within 30 days in the original packaging. Custom orders cannot be returned once the individual items are mixed. Bioengineering order cancellations are subject to our approval and assessed a 25% cancellation fee. Bioengineering orders are non-returnable.

Prices are subject to change without notice. Please call for pricing and availability.





YOUR PARTNER IN growth



Calvin & Marcia Ernst

Dear Friends,

In 1962, I planted my first five acres of crownvetch on my parent's farm for the steep price of \$200 per acre (\$2,000 per acre today), securing my first sale two years later. I was an eager college student who saw an opportunity and decided to take a risk. It certainly wasn't a move that paid off immediately, but it was one that embedded the seed of my adventurous agricultural career and lifelong commitment to conservation.

Today we are celebrating our 60th year, and what an honor that is. I could not be prouder of where we have come from and what we have grown into. Our increasingly expanding offerings increase the food supply and provide a lively habitat for bees, birds, and butterflies, as well as large and small mammals that enrich life for this generation and the next. We have also made exciting advancements in improving efficiency and increasing quality. The creation of custom software has allowed us to monitor and manage our crops and inventory by cutting time, not corners. Other changes, specifically in native seed production, include software and equipment upgrades that ensure we consistently offer the highest quality seed in the market.

The Ernst team has grown, along with our footprint, to 80+ members who oversee 200 species and manage 10,000 acres of fertile land. We have worked hard to cultivate a culture supportive of curiosity, encouraging experimentation, continued learning, and teamwork. It is a privilege to share every day with this talented, energetic, nature-loving group.

Looking ahead, I could not be more excited for the future or more eager to get up every morning and dig in. Thank you, our valued customers, for your continued commitment to conservation and unmatched loyalty.

Sincerely,

Calvin L. Ernst
Founder & President



OUR ROOTS

run deep



This year marks our sixth decade in the field and a lifelong commitment to conservation, but our story started with a curious boy.

Our founder and visionary, Calvin Ernst, was born and raised a stone's throw from today's sprawling agribusiness. At age 14, he obtained a Pennsylvania nursery license and began experimenting on his parent's farm. During his senior year of high school, he grew the record corn crop in Crawford County with guidance from his 4-H leader and the local Penn State Cooperative Extension office.

1960-69

As a junior at Penn State, Calvin, his brother Luther, and a few friends researched crownvetch for the Pennsylvania Department of Transportation (PennDOT). Learning of PennDOT's need and the lack of growers, Calvin and his brother scrape together \$1,000 to purchase seed and convince their father, Ted, to plant it on five acres of their family's farm.

Calvin and Marcia met and eventually married in 1963, forging their partnership as husband and wife and valued business partners.

Calvin and Luther planted an additional 60 acres of crownvetch on leased land. Their first crop was sold to Stanford Seed and, soon after, they hired their first full-time employee.

Development of the U.S. Interstate Highway System introduces I-79, I-80, I-90, and I-95 to the American landscape, accelerating the demand for erosion control and vegetation restoration.

The Company purchased a 180-acre property (the home of Ernst Seeds today) and joined forces with Multi-Systems, Inc., to purchase a 640-acre farm in Nebraska dedicated solely to growing irrigated crownvetch.

1970-79

The price of crownvetch collapses, and the partnership with Multi-Systems, Inc., is severed, leaving Ernst with a \$12,000 debt.

Calvin and Marcia secure financing for 500+ acres in Crawford County and expand by growing crownvetch as well as potted crownvetch plants, allowing them to pay off their debt.

1980-89

The Company continues to expand until the 22% interest rate hike, which caused the value of crownvetch to be less than the cost of harvesting it. Calvin pivots, planting large volumes of no-till corn in the existing crownvetch fields, creating high corn yields due to the legume's excellent nitrogen-fixing properties.

Ernst hires its first full-time IT employee to manage inventory, sales, and marketing activities, a progressive characteristic of the Company's professional and business-minded approach to agriculture.

The Company plants its first foundation native seed, *Panicum virgatum* (switchgrass) and *Panicum clandestinum* (Deertongue), and harvests native seed from surrounding wetlands in Crawford County.

Thoughts on non-natives evolve nationally, and the environmentalists blacklist some aggressive and invasive species. Calvin shifts his business strategy toward the diversity of natives, moving acreage from grain and crownvetch to native forbs and grasses.

1990-99

At the invitation of Rodale Institute, Calvin and Marcia, with their conservation experience, travel to eastern Europe to study diverse agricultural practices. Later, Calvin travels with the USDA to China to promote inter-cropping perennial legumes with annual grain crops to reduce crop inputs and erosion.

The Company thrives, harvesting and growing native wetland seeds for the wetland mitigation market and continues collaborating with federal agencies, highway departments, and land developers to establish wetlands and meadows using native seeds.

Ernst hires a full-time horticulturist to identify useful native species and develop production strategies for seed production.

Calvin and Marcia's children, Andy, Michael, and Robin, become involved in the Company, expanding native seed production and sales.

A fire destroys the Company's seed conditioning facility and several crops. Employees and neighbors rally, bringing tools to the farm to help the rebuild.

2000-09

Calvin utilizes vacuum harvesting technology on wiregrass in central Florida, and the Ernst family purchases a farm in Live Oak, Florida for field production. Due to decreased demand and production difficulties, the venture was not as fruitful as hoped. Still, it yielded a knowledge of Southeast native plants for reclamation and restoration.

Ernst leases 5,000 acres of farmland in Crawford County, plants switchgrass for seed and biomass production, and builds a state-of-the-art facility to process warm season grasses. Michael and a small group of employees design and construct a pellet plant that uses only grass. Today it operates as a separate entity, Ernst Biomass, LLC.

Andy and Michael continue in day-to-day operations at Ernst Seeds and Ernst Biomass, while Robin operates Meadville Land Service, Inc., and Monarch Vegetation Service.

2010-19

The Ernst Team consists of 80+ full-time employees, including field production crews, sales and marketing staff, researchers, seed conditioning specialists, finance, purchasing, and inventory management.

**2020-
PRESENT**

Ernst's owned acreage has doubled in the last ten years, and sales have increased by 50% in the last ten years. Healthy expansion continues with new drying and storage bins, equipment storage buildings, and a state-of-the-art spray bay.

As 2024 marks our 60th year in business, it will be the beginning of a new decade of growth and renewed excitement for decades to come.

MEET THE TEAM

At Ernst Seeds, we rise early, enjoy digging into a good challenge, and welcome any opportunity to share our knowledge from the field. Our highly skilled, expertly trained, and wildly innovative team runs 80+ members deep, with many celebrating over two decades at the farm. Needless to say, we take great pride in the work we do, the seeds we

offer, and the customers we serve. If you've got a question about growing, reach out to a pro today.

Got a question and are not sure where to start? Call (800) 873-3321, Monday through Friday, 8 a.m.–5 p.m. EST or e-mail us at sales@ernstseed.com.



Calvin Ernst
President



Marcia Ernst
Partner



Andy Ernst
Vice President



Michael Ernst
Vice President



Robin Ernst
Partner



Dan Arnett
Biomass Manager



Steph Breckenridge
*Operations & Inventory
Manager*



Paula Dithrich
*Senior Accounting
Manager*



Mark Fiely
Horticulturist



Cheri Haines
*Human Resources
Manager*



Kathy Haven
Executive Assistant



Nikki Hindle
Sales Manager



Kevin Jamison
Agronomist



Greg Kedzierski
*Plant Material
Specialist*



Need help after hours?

Check out our **Frequently Asked Questions** Section online if you need help after hours or over the weekend.



Sales & Order Processing



Warehouse & Shipping



Planting & Harvesting



Greenhouse



Summer Crew



Bioengineering



Seed Conditioning



Biomass



Accounting & Finance



Marketing

LET'S GET GROWING

DEFINING NATIVE VS. NATURALIZED SEEDS

NATIVE

Species that existed locally prior to European settlement.

NATURALIZED

Species not native to a certain area that grow, reproduce, and maintain themselves without interference.

ECOTYPE

Native species found in a defined area, state, or region.

VARIETY

A subdivision of native or naturalized species having one or more distinct, consistent, though often inconspicuous, traits.

WHY STOCK MIX FORMULATIONS CHANGE

Stock seed mix formulas may vary within a year or between years. Each mix is created with a particular guiding philosophy. As new species become available, they will be added to enhance the performance of the mixes.

Occasionally, a species may not be available due to crop failure or high sales, resulting in a reformulation using the remaining species in the mix. To adapt to these variations in our formulations, we recommend using the phrase "Ernst Mix (X) as currently formulated" when writing specifications.



ERNMX-140 Partially Shaded Area Roadside Mix



ERNMX-168 Northeast Annual & Perennial Wildflower Mix

UNDERSTANDING PURE LIVE SEED (PLS)

The Pure Live Seed (PLS) standard was developed to aid in determining the appropriate amount of bulk seed to be applied.

PLS refers to the percentage of live seed by weight in a seed lot having the potential to

develop into a seedling. Live seed refers to the germination percent, dormant seed, and hard seed. The remaining seed in the lot contains inert matter, other crops, and/or weed seeds and is not viable.

INDEX

Pure Seed – Viable and non-viable seed of the desired species.

% Purity – Percentage of pure seed by weight.

% Germination – Percent of seed by weight having the potential to germinate relatively quickly.

% Hard Seed – Percent of viable seed having a hard seed coat that can take longer to break and germinate.

% Dormant Seed – Percent of viable seed taking longer to germinate, other than hard seed.

% Total Germination – Percent of seed by weight composed of hard seed, dormant seed, and germination percentage having the potential to produce a seedling.

% Nonviable Seed – Percent of seed that will not germinate.

% Inert Matter – Material other than seed.

Viable Seed – Pure seed having the potential to produce a seedling.

Weed Seed – Seeds or other reproductive parts of species recognized by law as weeds.

Other Crop Seed – Seeds that are not weeds or pure seed.

HOW TO CALCULATE PLS

Percent Total Germination = (Germination + Hard Seed + Dormant Seed)

70 + 15 + 5 = 90% Total Germination

Use that figure and the purity percentage to calculate the PLS percentage.

Multiply Total Germination by Purity / 100 = PLS %
(95% X 90%) / 100 = 85.5% of Pure Live Seed


What Does This Mean?

85.5% of this seed lot by weight has the potential to grow. If 10 pounds of pure live seed is needed on a site, the quantity of bulk seed needed is determined by:


100 / PLS = Pounds of bulk seed needed to produce 1 lb of PLS

100 / 85.5 = 1.16 lbs of bulk seed X 10 lbs = 11.6 lbs of bulk seed

PURITY TEST



GERMINATION TEST



EXPECTANTIONS OF NATIVES SPECIES



EXPECTATIONS OF NATIVE SPECIES

GERMINATION & GROWTH (all the following assume adequate light, adequate soil temperature, appropriate moisture, and good seed-to-soil contact):

In general, annual species have less seed dormancy than biennials and biennials have less than perennials. Seed dormancy is nature's hedge against unfavorable conditions during a plant's life cycle. Unfavorable conditions can be, but are not limited to, late spring frost or drought. Dormant seeds germinate when favorable conditions are present.

ANNUALS

Most species germinate, flower, and set seed by the end of the first full growing season. Germination of an individual species is likely to be high.

BIENNIALS

Most species germinate, with some plants within a species' population flowering and setting seed in the first full growing season. The bulk of the plants will flower and set seed in the second growing season. Germination of an individual species is likely to be lower than that of an annual due to seed dormancy.

PERENNIALS

1. Warm Season Grasses: Germination will occur in the spring when moisture conditions are appropriate and soil temperatures exceed 55°F (12°C) at a 3" depth. Best germination occurs when soil temperatures are much higher.

Most of these species do not require cold, wet stratification to produce an adequate stand. Two exceptions are *Tripsacum dactyloides* (Eastern Gamagrass), which requires 14-60 days of stratification, and *Chasmanthium latifolium* (River Oats), which requires 60 days of stratification for northern genotypes. Stratification is the process by which seed is exposed to cool, moist conditions.

While cold, wet stratification is not necessary in most cases to produce an adequate stand, 20%-50% of the seed may be dormant. Most seedlings that emerge will be growing by the end of the second full growing season.

Greatest growth of these species occurs when air temperatures are 75°F-95°F (24°C-35°C). Most of the growth is in root development the first season. Very few (<5%) plants within a species may flower and set seed in the first growing season. Maximum plant development may take two years or longer.



New growth of Eastern Gamagrass (*Tripsacum dactyloides*) in a Florida field in the spring.



Eastern Gamagrass (*Tripsacum dactyloides*)

2. Cool Season Grasses: Some species will germinate when temperatures are a little higher than 40°F (4°C) while others will require warmer temperatures. They may germinate in the fall or spring. Adequate stands of most species do not require stratification; however, 50% of the seed may remain dormant without stratification. Most seedlings that emerge will be growing by the end of the second full growing season.

Greatest growth occurs when temperatures are 65°F-85°F (18°C-29°C). With adequate moisture and nutrients, some flowering and seed set may occur in the first growing season.

3. Some sedges (*Carex albolutescens*, *scoparia*, *vulpinoidea*), **rushes** (*Juncus effusus*, *tenuis*), and **bulrushes** (*Scirpus atrovirens*, *cyperinus*, *expansus*, *polyphyllus*) have a very high seed count per pound of seed. When planted in the spring, a substantial number of seedlings may be produced by these species in the first growing season. These seedlings may represent 5% or less of the total seeds present. Flowering and seed production will occur one to two growing seasons after an individual seedling has germinated. Maximum germination will take at least two years due to seed dormancy. Sedges and bulrushes are recognizable by the arrangement of any three successive leaves in a pattern resembling the spokes in the Mercedes™ symbol. *Juncus spp.* will have round stems that originate at a common point near or on top of the soil.

4. Some sedges (*Carex comosa*, *crinita*, *frankii*, *glaucescens*, *grayi*, *intumescens*, *lupulina*, *lurida*, *squarrosa*, *stricta*) and **bulrushes** (*Scirpus validus*) have a high level of seed



Virginia Wildrye (*Elymus virginicus*)

dormancy and may not have consequential germination without stratification.

Most seedlings will emerge in the first and second growing seasons after stratification (artificially or naturally). Plants will flower and set seed one to three years after they germinate. *Carex spp.* in this group may be recognized as described above for other *Carex spp.* *Scirpus spp.* in this group have round or triangular stems arising from a point often below the soil surface. The stems are typically larger than those of *Juncus spp.*

5. For most broadleaf species, some germination will occur in the first year without stratification (artificial or natural). A high percentage of species and seeds within the species are likely to germinate in the first growing season following the first winter *in situ* (on-site). Most of the seeds that germinate will have done so by the end of the growing season following stratification. Following germination, blooms may occur in the first growing season: *Heliopsis helianthoides* (Oxeye Sunflower); second growing season: *Rudbeckia triloba* (Brown-eyed Susan), *Aster spp.*, *Monarda spp.*, *Penstemon spp.*, *Solidago spp.*; after three to five growing seasons: *Liatris spp.*; or, not until the seventh growing season: *Baptisia tinctoria* (Yellow False Indigo). The number of blooming years depends on soil fertility, available moisture, and growing season temperatures. It may be shorter for a given species the further south one is located.

6. Seed dormancy in perennial species is affected by latitude of ecotype origin. In greenhouse studies, we have found that



Nodding Sedge (*Carex crinita*)



Blooms may occur in the first growing season with Oxeye Sunflower (*Heliopsis helianthoides*), while other broadleaf species may take numerous growing seasons to bloom.

northern ecotypes (PA, OH, NY, NJ) typically require more weeks of cold, wet stratification than southern ecotypes (FL, GA, NC, SC) of the same species.

Most of our native seed mixes are composed of perennial species. Mixes dominated by perennial species have the potential to last more than a decade if properly maintained. For all mixes, a site must be kept free from invasive species or aggressive weeds. Mixes of herbaceous species with no tree, shrub, or vine components in the formula must be kept free from the encroachment of woody or vine species with controlled burning, mowing, or spot spraying.

The natural communities we create with native seed mixes are dynamic. Annuals, biennials, and short-lived perennials may be widely present in the landscape in the first three growing seasons, but non-existent or present in small pockets by the fifth growing season. Over time, colonies of some long-lived perennials will grow larger in area and species composition will change in response to annual rainfall variations.

It is not unusual for those new to planting meadows to be nervous about a mix's performance during its establishment year. Typically, customers need confirmation that the desirable species are growing. Fortunately, our ability to assess a situation is assisted by a small set of species that generally germinate very well.

For wetland meadows, some common early emerging species include: *Asclepias incarnata* (Swamp Milkweed), *Eupatorium perfoliatum* (Boneset) and *Carex spp.* For upland meadows, some common early emerging species include: *Chamaecrista fasciculata* (Partridge Pea), *Elymus virginicus* (Virginia Wildrye), *Helianthus angustifolius* (Narrowleaf Sunflower), *Heliopsis helianthoides* (Oxeye Sunflower), *Monarda fistulosa* (Wild Bergamot), *Penstemon digitalis* (Tall White Beardtongue), and *Rudbeckia hirta* (Blackeyed Susan). Seedling images of many of these species are available on our website.

DISCLAIMER: The information in this review of practices is the result of 60 years' experience in seed production. Ernst Seeds has been supplying seeds and consulting in the reseeding of tens of thousands of acres of roadsides, surface-mined lands, conservation, and restoration sites in eastern North America, as well as growing and supplying seed and consulting in the planting of hundreds of thousands of acres of CRP/CREP-related areas for erosion control and wildlife habitat.

All these practices are opinion only and our best advice as a result of these experiences. These recommendations are for individual consideration and do not cover all the conditions that will be encountered in the field.

Ernst Seeds is not responsible for conditions that will be encountered in individual situations. The use of brand names does not represent our endorsement of a specific product; rather, it represents our experience only and has not necessarily been replicated in peer-reviewed research. The use of chemical pest control agents is subject to manufacturers' instructions and labeling, as well as federal, state, and local regulations.

ESTABLISHMENT GUIDE



ERNMX-105 Mesic to Dry Native Pollinator Mix buffer planting.

ESTABLISHMENT GUIDE INTRODUCTION

In eastern North America, there is a wide variety of native vegetation to replicate. Most planting objectives fall into the following categories:

- Soil erosion control & soil stabilization on slopes and along waterways
- Beautification & enhancement of landscapes
- Biodiversity & wildlife habitat enhancement and restoration
- Bioremediation to correct environmental disturbances
- Historical, cultural & ecological restoration
- Habitat for honeybees & native pollinators (butterflies, bumblebees, etc.)
- Native species for renewable biomass production

Using native plants saves time and money while improving ecological function. Reduced water, chemical, fertilizer, and maintenance needs make them a sustainable and environmentally sound choice for virtually all scenarios. Select a mix of species that creates the desired outcome for the project. Goals should be compatible with site conditions that cannot be altered. Native plant communities can be selected to meet nearly all site conditions.

Please review the appropriate section(s) below for information regarding seed mix selection and seeding methods. Matching the functional goals of the site and site conditions to the appropriate seed mix will lead to greater project success. The stock seed mixes noted in each section represent a mere sampling of our complete list of mixes. A more comprehensive list may be found at www.ernstseed.com or by contacting a member of our sales team. Mixes can also be customized to specific needs as well as those of a site and ecological region.

PLANTING GUIDES

UPLAND & MEADOW SITES	23
WILDLIFE HABITAT & FOOD PLOT SITES	27
POLLINATOR-FRIENDLY SITES	30
RIPARIAN SITES	33
DISTURBED SITES & STEEP SLOPES	36
UTILITY RIGHT-OF-WAY SITES	40
WET MEADOW & WETLAND SITES	42
PARTIALLY SHADED SITES	46
STORMWATER MANAGEMENT SITES	50
SOUTHEASTERN U.S. SITES	54
BIOMASS PRODUCTION SITES	68

FALL VS. SPRING SEEDING

Traditionally, seeding is thought of as a spring activity, but it can also occur during the dormant season. Fall seeding works well for restoration projects completed in the summer.

While there are some noteworthy advantages to fall seeding, seeding in either spring or fall will produce good results. In drought-prone regions, seeding should be timed to take advantage of the available moisture in the area.

FALL OR "DORMANT" SEEDING

- ▼ Fall seeding imitates natural reseeding. Dormant seeding can take place when soils are dry enough to work.
- ▼ Good seed-to-soil contact occurs through precipitation and the freeze-thaw cycle.
- ▼ Natural stratification and scarification occur; natural changes within the seed or to the seed coat during the winter enhance germination in the spring.
- ▼ Mulching is an important element of dormant seeding to protect the soil.
- ▼ Some seed may be lost to decay and wildlife consumption during the winter.
- ▼ Establishment may be hindered by growth of winter annuals in the fall.

FROST SEEDING

- ▼ Frost seeding is the act of broadcasting seed onto or drilling into frozen soil.
- ▼ Good seed-to-soil contact occurs through the freeze-thaw cycle.
- ▼ Natural stratification and scarification occur; natural changes within the seed or to the seed coat during the winter enhance germination in the spring.
- ▼ Mulching may be done.
- ▼ Some seed may be lost to decay and wildlife consumption during the winter.
- ▼ Establishment may be hindered by growth of winter annuals in the fall.

SPRING SEEDING

- ▼ Cool season species germinate soon after seeding.
- ▼ Germination of warm season species generally occurs within three weeks of the soil temperature reaching 55°F (13°C).
- ▼ Seed loss due to decay and wildlife consumption is minimized.
- ▼ Seed-to-soil contact should be accomplished by working the seed into the soil 1/4"-1/2" deep.
- ▼ Seeding may be delayed until weed control is applied to improve establishment.
- ▼ Irrigation during dry weather periods is necessary for proper germination.
- ▼ Light mulching is an important element of seeding to protect both the seed and soil and retain moisture.
- ▼ When planting native grasses for biomass, seeding generally takes place during the spring when soil temperatures at a 3" depth are near 55°F and rising.



SEEDING METHODS



DRILL SEEDING

Drill seeding is a mechanical means of creating furrows (openings) in the soil surface and metering seed in at a uniform rate. A drill seeder is practical for seeding multiple acres in larger areas.

Conventional drills can work in tilled and partly tilled soil. No-till drills are designed to work in soil that has not been tilled. They have heavy openers that cut through vegetation and sod to make a furrow for seed placement. With the proper adjustment, a no-till drill can work in tilled soil. It has discs that aid in loosening the soil. All drills should be equipped with a closing or packing wheel that follows seed placement.

The goal of drill seeding is to achieve uniform seed distribution over the site with seed placement at the correct depth (1/4"-1/2") and good seed-to-soil contact. Calibrating a drill or broadcast seeder depends on seed bulk density and required application rates. Manufacturers provide manuals with charts to guide seeding rate calibration. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area, then make any necessary adjustments. Most traditional seed drills are designed to handle seeds with high bulk densities, such as oats and wheat. Some drills may have a small seed box able to plant small seeds, such as alfalfa, clover, and switchgrass.

Many native and naturalized species are fluffy and will not readily flow through a traditional seed drill. Examples of fluffy seed include little bluestem, big bluestem, and indiangrass. With the aid of a bulking agent, some fluffy seeds may be planted through the large seed box of a traditional drill. Bulking agents include kitty litter, dry sawdust, vermiculite, or rice hulls. Test with a small amount of seed. Native seed drills, such as Truax, have specialized seed

boxes that are effective for planting fluffy seed. When seed will not readily flow through a native seed drill's fluffy seed box, a bulking agent may be needed.

HAND SEEDING

Hand seeding is the casting of seed onto the soil. Hand seeding is used on small plots or difficult terrain where seeding with machinery is not an option. The goal is to achieve an even distribution of seed over the site. This can be accomplished by spreading half of the seed in one pass and the balance in a perpendicular pass. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area. To know how wide to make your passes, check the width of seed distribution.

If possible, a light raking to a depth of 1/4" and/or firming with a lawn or Brillion-type roller is recommended to achieve good seed-to-soil contact. Cover with straw mulch at 70 lb per 1,000 sq ft or hydromulch at 34 lb per 1,000 sq ft.

When the volume of seed to be applied is small (less than 50 lb per acre), a bulking agent may be helpful to provide the volume necessary to get uniform application. Such bulking agents include kitty litter, dry sawdust, vermiculite, or rice hulls.

BROADCAST SEEDING

A broadcast seeder consists of a hopper with an adjustable door that regulates seed flow onto a spinner. Some broadcast seeders have an agitator that aids with seed flow in the hopper. Broadcast seeders are commonly used to spread seed, fertilizer, lime, and other granular products. The goal is to achieve an even distribution of seed over the site. To ensure uniform application of seed, conduct a test run over a small area using the appropriate amount of seed for that area. To know how wide to make the passes, check the width of seed



distribution from the spreader. The settings can then be adjusted as needed. To achieve better distribution, spread half of the seed in one pass and the balance in a perpendicular pass. We recommend refilling the hopper when it is 1/3 full rather than letting it empty out. Follow up by tracking or firming the seed into the soil with a lawn or Brillion-type roller to achieve good seed-to-soil contact. Do not roll or track the seed if the soil is wet. Cover with straw mulch at 70 lb per 1,000 sq ft or hydromulch at 34 lb per 1,000 sq ft.

Many native seeds are fluffy and will not uniformly flow through a broadcast seeder. To enhance the flow, mix the seed with a bulking agent of similar density. Dry sawdust, vermiculite, or rice hulls are some options. An agitator in the hopper may be required in these circumstances. We recommend a minimum rate of 50 lb per acre of seed and bulking agent.



A bulking agent can also be helpful if you are planting small quantities of seed. It provides the volume necessary to get uniform application. For fine seeds, kitty litter is a more appropriate bulking agent.

CULTIPACKING

A cultipacker is an excellent way of covering the seed with a minimum amount of soil to ensure proper seed-to-soil contact. It resembles a large rolling pin with evenly spaced ridges and dimples. The primary functions of a cultipacker are to break up clods, remove excess air spaces from loose soil, and smooth the soil. The heavy-duty



smooth, spoke, or crowfoot rollers provide clod-breaking and smoothing capabilities. As with any tillage, it is important not to overwork the soil or work it when it is too wet.



HYDROSEEDING

A hydroseeder combines water, seed, fertilizer and, sometimes, hydromulch into a mix that is pumped through a nozzle and sprayed uniformly over the area to be seeded. Hydroseeders can distribute this mix at 150' or more, allowing for the ability to seed terrain that may not be accessible with other seeding methods, such as steep slopes, roadside cuts, or sites that are too wet. Using hydromulch aids in seed placement and reduces erosion on slopes. Depending on site conditions, use of erosion control blankets or straw mulch may be needed to cover the seed. Many native seeds should be broadcast with 500 lb per acre of mulch as a marker. Do not exceed this amount as native seeds may die if suspended in the mulch with little or no seed-to-soil contact. The balance of the hydromulch, often 1,000 lb per acre, may be applied on top in a secondary application.



STRAW MULCHING

A straw-mulch blower distributes mulch over a seeded area. It has a slide (or chute) in which to feed the mulch, chopper blades to break up the mulch, and a blower to spread the mulch over large areas. Straw mulch may be spread by hand in smaller areas. It is important to use weed-free straw from small grains, such as oats or grain rye, to minimize potential weed issues.

TOOLS FOR PREPARATION



MINIMUM-TILL EQUIPMENT

Minimum-till equipment incorporates a portion of the surface vegetation into the soil and levels uneven surfaces. One of the most common tools is a disc which cuts through vegetation, sod, or hard soil and partially turns or tills it into the soil. Similar equipment that turns part of the vegetative residue into the soil is known as Aerway® or Turbo® Till.



TRACKING

Tracking is the use of a crawler or rubber-tired tractor to make depressions and firm loose soil after construction or tilling. Tracks should be oriented perpendicular to the slope of a site. The depressions from tracking help to reduce erosion and retain seed and moisture. The firm, but not compacted, seedbed will not dry out as quickly as loose soil.



CHISEL PLOW

A chisel plow is a minimum-till plow because it does not dislodge or turn over the entire soil profile the way a moldboard plow does. Chisel plowing is primarily used for breaking up hardpan soil or loosening compacted soil while leaving a high percentage of debris on top. A chisel plow can be adjusted to till shallow or deep and typically has C-shaped shanks mounted on dual coil springs, and the frame, shanks, and springs are of sufficient weight, size, and strength to provide a cutting depth of 8"-12". To make the soil smooth enough for planting after the use of a chisel plow, use a disc harrow, tandem disc harrow, or offset disc harrow of sufficient weight and size to provide a cutting depth of 6"-8".



ROTOTILLER

A rototiller is used to pulverize the soil with rotating blades and incorporate soil amendments and surface vegetation. Most units till up to 6" deep.

TOOLS FOR MAINTENANCE



ROTARY MOWER

Heavy vegetation on under-utilized fields is difficult to mow with a discbine or sickle bar mower. Heavy-duty rotary mowers can be utilized as brush hogs to tame heavy grass and light brush, such as multiflora rose, honeysuckle, and small tree seedlings.



SPRAYER


Sprayers come in various sizes and styles, including common hand-held units like the one shown here. These are often preferred for carefully targeted spraying of unwanted or invasive vegetation. Larger areas may be effectively sprayed using tractor or ATV-drawn tank units.

Use of herbicides to control undesirable vegetation can be an important part of an integrated pest management (IPM) program when applied according to the manufacturer's label. Prior to using any herbicide, read the label for safe handling and application information. Many herbicides are only available to licensed applicators. When these are needed, employ a licensed professional.



DISCBINE MOWER

A discbine mower is a hay-harvesting machine with high-speed rotary discs that mow biomass for baling and assemble the material into a windrow.



Mowing during the growing season should not be necessary after the establishment year unless it is in lieu of herbicides to control weeds. Mowing height should be no lower than 8".

To prevent succession of woody species in an established meadow, an important aspect of a maintenance program is an early spring mowing that is close to the ground (2"). Mowing should take place every one to three years, in late winter or early spring, and shortly before spring nesting season. Spring mowing will leave cover and food for wildlife through the winter without disrupting the nesting of grassland birds.



UPLAND & MEADOW SITES

FAST FACTS

Upland & Meadow Sites are characterized as being dry most of the year. Soils on these sites are well-drained and may consist of sandy clay, sandy loam, loam, or shale. The topsoil layer may be thin and subject to drought. If the site is sandy, refer to the Southeastern U.S. Sites Establishment Guide, p. 54.

Upland examples: Naturally rocky soil subject to erosion or steep road cuts. Meadow examples: abandoned farm fields, lawns, vacant land, or roadsides.

SITE PREPARATION

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate time interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedlings from germinating.


Competition from invasive or undesirable vegetation is the most limiting factor for a successful upland meadow establishment. Prior to planting, all such vegetation must be fully controlled. It may take a full growing season or more to control rhizomatous species, such as mugwort, Canada thistle, poison ivy, or johnsongrass. If these species are not fully controlled prior to planting, they will overrun your planting. Typical control strategies include repeated tilling, smothering with black plastic, or herbicides. While any of these methods may control existing weeds, they will not kill all weed seeds lying dormant in the soil. Seeds of such species as velvetleaf and pokeweed may germinate many decades after the species last flowered on the site.

When using the tillage strategy, a site is disc harrowed every two to four weeks for one to two months. The underlying principle of this process is that the root system of perennial

species will be worn out to the point of killing the species. In addition, tillage will stimulate germination of some dormant weed seeds that will be killed with subsequent tillage. Planting should not take place until perennial species are completely killed.

Black plastic may also be used to kill weeds. It may be laid across tilled or untilled soil and anchored down by burying the edges in soil or laying boards or bricks across the surface. This strategy should be utilized in a growing season when the intent is to fall plant in the same year or spring plant the following year.

Use of an approved herbicide, such as glyphosate (Roundup® or Rodeo®), by a licensed spray technician is the most common and least time-intensive method for controlling existing vegetation. Since herbicides are most effective on actively growing plant tissues, they are very effective on new spring growth. Spraying should begin when growth reaches 6". A follow-up application one to two weeks later will address skips or persistent species. If substantial plant tissue remains on the surface following a full kill by herbicides, close mowing, tillage, or burning may be necessary to achieve good seed-to-soil contact.



ERNMX-179 Butterfly & Hummingbird Garden Mix meadow planting in Hershey, Pennsylvania.



ERNMX-179 Butterfly & Hummingbird Garden Mix meadow planting in Hershey, Pennsylvania.



An upland meadow using ERNMX-179 Butterfly & Hummingbird Garden Mix. Credit: Paul Stead

GROWING SEASON MAINTENANCE

MAINTENANCE

Problem weeds should be hand pulled in annual wildflower mixes or annual and perennial wildflower mixes. For all other mixes, see below.

FIRST GROWING SEASON

Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed. Trimming should cease by mid-September.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Sonora™, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

SECOND & SUBSEQUENT GROWING SEASONS

Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any remaining material from the previous year close to the ground (approximately 2"). This will allow the soil to warm more quickly, stimulating emergence and growth of native seedlings and reducing the likelihood of shrub invasion.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Sonora™, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

SPECIAL CIRCUMSTANCES

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September. To prevent weed reinfestation, use of an appropriate selective herbicide in conjunction with a seed mix tolerant of that herbicide may be necessary.



HABITAT

Upland to mesic sites, typically in full sun for at least half the day with good air circulation.



FERTILITY

Natural fertility is usually adequate; fertilizer and lime are not needed. Check soil pH and select appropriate species.



SEEDING METHOD

Hand seed, broadcast seed, hydroseed, or drill seed.



A meadow installation at a Princeton, New Jersey residence. Credit: OPEN Landscape Architecture

UPLAND & MEADOW SITES SEED MIXES

ERNMX-105	Mesic to Dry Native Pollinator Mix
ERNMX-110	Ernst Native Biomass Mix for Strip Mines & Gas Production Sites
ERNMX-111	Ernst Native Habitat for Strip Mines Mix
ERNMX-115	Biodiverse Polyculture Mix for Biomass Production & Wildlife Habitat
ERNMX-117	Warm Season Grass Mix
ERNMX-123	Native Upland Wildlife Forage & Cover Meadow Mix
ERNMX-125	Mesic to Dry Native Pollinator Mix without Grasses
ERNMX-153	Showy Northeast Native Wildflower & Grass Mix
ERNMX-153-1	Showy Northeast Native Wildflower Mix
ERNMX-156	Low-Growing Wildflower & Grass Mix
ERNMX-166	Plateau-Tolerant Wildflower & Grass Mix
ERNMX-166-1	Plateau-Tolerant Wildflower Mix
ERNMX-167	Annual Wildflower Mix
ERNMX-168	Northeast Annual & Perennial Wildflower Mix
ERNMX-169	Southeast Annual & Perennial Wildflower Mix
ERNMX-170	Annual & Perennial Wildlife Food Plot Mix
ERNMX-171	Multi-Purpose/Multi-Year Wildlife Food & Shelter Mix
ERNMX-172	Maryland Upland Mix
ERNMX-173	Eastern Native Habitat & CREP Mix
ERNMX-174	Virginia Gentleman's Mix
ERNMX-177	Eastern Ecotype Native Grass Mix
ERNMX-179	Butterfly & Hummingbird Garden Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.

Seeds that feed!

These mixes are used in full sun with well-drained soils and provide food and/or cover for wildlife. Meadow and wildflower mixes provide food for insects, including native pollinators.

WILDLIFE HABITAT & FOOD PLOTS SITES

FAST FACTS

Wildlife Habitat and Food Plot Sites are generally small clearings in wooded areas or farmland. While planting dates differ between annual and perennial mixes, most are designed for spring or fall planting. Examples: Woodland openings, remote areas of large residential lots, abandoned fields, or timber harvest and loading areas.



SITE PREPARATION

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedling germination.

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Roundup® or Rodeo®), or by tilling the weeds into the soil. Hand pull or spot spray problem weeds. Perennial weeds not addressed before establishment will be difficult to remove later. Good pre-seeding weed control may require repeated tilling or spraying two applications of glyphosate at least two weeks apart.

GROWING SEASON MAINTENANCE

It may be necessary to mow some mixes to 4"-6" in order to keep plants young and tender. Annual food plot mixes must be planted yearly.



HABITAT

Well-drained or moderately well-drained sites in woodland openings (ideally near water sources); typically in full sun for at least half the day; upland species may be planted.



FERTILITY

Check soil pH and fertility; adding lime can improve the nutritional value of vegetation beneficial for wildlife. If a soil test has not been obtained, a starter fertilizer, such as 200 lb per acre of 16-16-16, should be applied. If uncertain about the soil pH, add 1,000-2,000 lb of lime per acre which will provide plants with essential nutrients without pushing up a lot of top growth. Lime and fertilizer may be incorporated into the soil using a tiller. After incorporating amendments, smooth the soil to develop a good seedbed. If broadcasting the seed, run a spring-tooth harrow or ATV over the site to incorporate the seed into the soil 1/4"-1/2" deep.



SEEDING METHOD

Hand seed, broadcast seed, or drill seed.




ERNMX-184 Fall Sweets Wildlife Mix. Credit: Elicia Winner.



A ringneck pheasant rooster takes flight in a hunting preserve using Ernst habitat mixes.



A pair of eastern wild turkey jakes in a spring food plot.



A mature whitetail deer standing on the edge of a marsh.

WILDLIFE HABITAT & FOOD PLOT SITES SEED MIXES

ERNMX-130	Wildlife Food Plot Mix
ERNMX-133	Keystone Deer & Turkey Habitat Mix
ERNMX-133-1	Keystone Big Buck Mix
ERNMX-170	Annual & Perennial Wildlife Food Plot Mix
ERNMX-171	Multi-Purpose/Multi-Year Wildlife Food & Shelter Mix
ERNMX-184	Fall Sweets Wildlife Mix
ERNMX-185	Spring Greens Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.



POLLINATOR- FRIENDLY SITES

FAST FACTS

Pollinator meadows may be used to attract honeybees and more than 4,000 species of native pollinators in North America, including bees, butterflies, hummingbirds, and even some flies. Increased attention to the plight of such pollinators as honeybees and monarch butterflies by the government and private sector has spurred an interest in developing pollinator-friendly habitats across the North American landscape. One of the highest examples was the 2015 White House announcement of the National Strategy to Promote the Health of Honeybees and Other Pollinators.



ABOUT POLLINATOR-FRIENDLY SITES

Almost any site not intended to be mowed repeatedly during the growing season may be designed to be pollinator friendly. Pollinator-friendly meadows are often thought of as being planted on flat sites in full sun, but they can also be planted at woodland edges. Pollinator-friendly species aid in soil stabilization on steep slopes and riparian areas. Storm basins and wetlands are more aesthetically pleasing when enhanced with pollinator-friendly species. Rights-of-way under utility transmission lines, above pipelines, and along roadways can be developed to an ecologically beneficial state when functional diversity and pollinator-friendly species are incorporated into seed mix design. When planted within a solar array, transpiration from native plants can reduce panel temperatures thereby increasing panel efficiency.

The primary energy source for most adult bees, butterflies, and other flower-loving pollinators is nectar. Pollen is essential for providing proteins and lipids to developing bee larvae while leaf tissue from specific host plant families is required for butterfly caterpillars. Most native bees are nectar generalists in that, though pollen specialists, they can consume nectar from many plant families. They are also pollen

specialists whose larvae require a specific ratio of proteins to lipids. The best sources of pollen for native bees as well as leaf tissue for native butterflies are the native plant species with which they have co-evolved.

While not native to the U.S., honeybees have evolved to be able to use pollen from a wide range of species. Like native bees, honeybees feed nectar and pollen to their larvae. They also need pollen to have a particular protein-to-lipid ratio that they get by collecting pollen from a variety of plant species.




Honeybee on a Cup Plant (*Silphium perfoliatum*).



Solar power generation sites can be sources for pollinator habitat.

Adult pollinators do not have the same dietary needs as when in the juvenile stage. To meet the dietary needs of a wide range of pollinators, it is important to know that some pollinator species are not active for the entire growing season. During the active period, food and nesting resources must be available. The availability of flowering shrubs or trees for pollen and/or

nectar before herbaceous species bloom in the spring is beneficial to some pollinator species. Continuity of bloom from as early in the season to as late in the season as possible is important. A minimum of three species should be in bloom in the spring, summer, and fall. For the benefit of monarchs, milkweeds should be planted.



Swallowtail Butterfly on Great Blue Lobelia (*Lobelia siphilitica*).

TO SUPPORT THE GREATEST DIVERSITY OF NATIVE POLLINATORS:

- Provide continuity of bloom from as early to as late in the season as possible.
- Minimum of three species blooming in spring, summer, and fall.
- Plant milkweeds for monarchs.



POLLINATOR-FRIENDLY SITES SEED MIXES

ERNMX-105	Mesic to Dry Native Pollinator Mix
ERNMX-125	Mesic to Dry Native Pollinator Mix without Grasses
ERNMX-153	Showy Northeast Native Wildflower & Grass Mix
ERNMX-153-1	Showy Northeast Native Wildflower Mix
ERNMX-155	Deer-Resistant Meadow Mix
ERNMX-157	Honeybee Forage Mix
ERNMX-179	Butterfly & Hummingbird Garden Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.

RIPARIAN SITES

A close-up photograph of a flowering plant, likely a species of milkweed, with clusters of small pink and white flowers. A bee is visible on the right side of the flower cluster. The background is a soft-focus green, suggesting a natural outdoor setting.

FAST FACTS

Riparian Sites are usually adjacent to rivers and waterways with soils often containing clay, high amounts of organic matter, and/or saturated sand. Examples: River and streambanks or damp floodplains of rivers and streams.



SITE PREPARATION

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Rodeo), triclopyr (Garlon® 3A), or a similar aquatic herbicide formulation, to control such undesirable vegetation as multiflora rose, honeysuckle, and woody species. CAUTION: Some persistent species, such as purple loosestrife, phragmites, Japanese knotweed, or reed canarygrass, may require multiple applications of glyphosate or triclopyr. Perennial weeds not addressed before establishment will be difficult to remove later. Before seeding, excess dead vegetation should be removed, turned under, or burned if conditions permit. Newly constructed riparian sites should be seeded as soon after construction as possible.



HABITAT

Varies from partial shade to full sun; subject to flooding; generally populated with riparian and floodplain species.



FERTILITY

Due to the potential for water contamination, lime and fertilizer are not recommended; however, we do recommend the addition of organic matter (straw, compost, mulch, leaf litter, etc.) when topsoil has been depleted or removed. Check soil pH and select species adapted to that pH.



SEEDING METHOD

Hand seed, broadcast seed, or hydroseed.



ERNMX-178 Riparian Buffer Mix along a small stream corridor.

Before



After



Bowman's Creek re-setting and rehabilitation project in northeastern Pennsylvania. Credit: Adam Nordfors

GROWING SEASON MAINTENANCE

FIRST GROWING SEASON

Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, or Garlon® 3A. Be vigilant in controlling invasive vines, such as bindweed, mile-a-minute, and Japanese hops. These are more easily pulled early than after two to three months of growth. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

SECOND & SUBSEQUENT GROWING SEASONS

Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any remaining material from the previous year close to the ground (approximately 2"). This will

allow the soil to warm more quickly, stimulating emergence and growth of native seedlings and reducing the likelihood of shrub invasion.

If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo® or Garlon® 3A. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September.


GENERAL MAINTENANCE

Grassy weeds or persistent perennials can re-establish in these soils. Monitor and control weeds by hand pulling or spot spraying.

RIPARIAN SITES SEED MIXES

ERNMX-154	Floodplain Mix
ERNMX-178	Riparian Buffer Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.





DISTURBED SITES & STEEP SLOPES

FAST FACTS

Disturbed Sites & Steep Slopes have various soil types and conditions typically distinguished by lower quality soils and a predisposition to runoff and erosion. Examples: Landfills, surface mines, road cuts, or construction sites.

SITE PREPARATION

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide. Perennial weeds not addressed before establishment will be difficult to remove later. Whenever possible, regrade the site to reduce slope and build diversions to reduce erosion and minimize seed loss.

For areas with slope greater than 3:1, final tracking should be perpendicular to the slope. The tracks will aid in reducing erosion and retaining seed and moisture.

Mulching with straw, hydromulch, or straw/coconut fiber mats is recommended on these sites to protect the seed from drying out or washing away. For areas steeper than 3:1, the use of erosion control blankets or flexible growth medium (e.g., Flexterra®) is recommended. When using erosion control blankets, be sure they are toed in at the top of the slope.



HABITAT

Various soils with exposed clay, sand, and rock outcropping without topsoil as a result of construction; generally populated with upland species.



FERTILITY

Typically low in fertility; therefore, adding topsoil or organic matter (compost) can be very beneficial. Check soil pH and select species adapted to that pH. Add lime and fertilizer as recommended by soil analysis. Incorporate amendments into the soil in a way that will leave the soil rough and minimize soil erosion and rapid runoff (e.g., tracking). If there is a weed problem, fertilizing is not recommended.



SEEDING METHOD

Hand seed, broadcast seed, hydroseed, or drill seed. For areas with slope less than 3:1, cover the seed 1/8"-1/4" deep by dragging with a spring-tooth harrow or firmly pressing the seed into the soil using a cultipacker, lawn roller, or ATV.



ERNMX-181 Native Steep Slope mix with Annual Ryegrass in Morgantown, West Virginia.



A one-year-old Big Bluestem (*Andropogon gerardii*) meadow at Fishkills Landfill on Staten Island, New York.



This wetland was constructed on a former mine site.

GROWING SEASON MAINTENANCE

FIRST GROWING SEASON

Post-planting maintenance will provide improved results if the ground is not too rough or steep. Whenever canopy height (overall vegetation) reaches 18"-24", use a brush hog mower or string trimmer to trim the meadow to 8". Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Sonora™, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

SECOND & SUBSEQUENT GROWING SEASONS

Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2") on sites that are not too rough or steep. This will allow the soil to warm more quickly, stimulating emergence and growth of native plants and reducing the likelihood of shrub invasion.

If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Sonora™, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September. However, vegetation allowed to grow without mowing provides more protection for wildlife and aids in erosion control.



A steep slope and retaining wall utilizing ERNMX-181 Native Steep Slope Mix with Annual Ryegrass at the Millcreek Mall in Lancaster, Pennsylvania.



Seedlings from a steep slope mix poking through an erosion control blanket.



DISTURBED SITES & STEEP SLOPES SEED MIXES

ERNMX-101	Non-Native Ernst Best Strip Mine & Gas Production Mix
ERNMX-102-1	Pipeline Mix with Switchgrass
ERNMX-103	Non-Native Good Value Mine Mix
ERNMX-104	Quick Erosion Control Cover Mix
ERNMX-109	Crownvetch Seeding Mix (Naturalized)
ERNMX-110	Ernst Native Biomass Mix for Strip Mines & Natural Gas Production Sites
ERNMX-111	Ernst Native Habitat Mix for Strip Mines
ERNMX-181	Native Steep Slope Mix with Annual Ryegrass



These mixes are good for controlling erosion and providing food and/or cover for wildlife.

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.

UTILITY RIGHTS- OFF-WAYS

FAST FACTS

Many utility companies and oil and gas operators recognize a unique opportunity to go a step beyond basic reclamation standards by using native species indigenous to the area being restored. In doing so, they are affirming the industry's commitment to environmental stewardship.



In the years after disturbance, affected areas can become biodiverse ecosystems with improved ecological function, greater wildlife populations, less erosion, and improved water and soil quality.

For example, a multiple-mile stretch of pipeline in the Marcellus and Utica shale plays may pass through wetlands, over steep mountain slopes, across rivers, and through agricultural areas and state game lands. These areas should be reclaimed with vegetation best matching the intended use of the land, the biodiversity that existed before disturbance, and with practices that best address such issues as erosion control, habitat fragmentation, and other environmental concerns.

We can design a biodiverse native seed mix to mitigate the environmental impact and aid in ensuring regulatory compliance. We routinely work with environmental departments, consulting engineers, and contractors seeding a project.

For Upland Meadow Sites Establishment Guide, see p. 23;
Disturbed Sites & Steep Slopes Establishment Guide, see p. 36;
Wetland Sites Establishment Guide, see p. 42;
Riparian Sites Establishment Guide, see p. 33.



This former well pad is bringing ecological value back to the landscape.



A pipeline seeded with a mix of native grasses, forbs, and legumes.

WET MEADOW & WETLAND SITES

FAST FACTS

Wet Meadow & Wetland Sites have soils made up of clay and high organic matter with high water tables or impervious layers that prevent drainage. These sites are wet most of the time. Wet Meadow Examples: Roadside ditches, retention basins that catch run-off water, pond areas, or wetland edges. Wetland Examples: Newly created wetlands and wetland restoration sites, retention basins with wetland functions, floodplains, pond edges, open water, or wet bioremediation sites.



SITE PREPARATION

Eradicate existing vegetation by having a licensed spray technician apply an approved herbicide, such as glyphosate (Rodeo), triclopyr (Garlon® 3A), or a similar aquatic herbicide formulation, to control such undesirable vegetation as multiflora rose, honeysuckle, and woody species. CAUTION: Some persistent species, such as purple loosestrife, phragmites, Japanese knotweed, or reed canarygrass, may require multiple applications of glyphosate or triclopyr. Perennial weeds not addressed before establishment will be more difficult to remove later. These sites are often too wet to till. Newly constructed wetlands, retention basins, and wet construction sites should be seeded as soon after construction as possible. Leaving the surface rough by creating mounds and kettles for an undulating microtopography can be very beneficial in obligate wetlands.



HABITAT

Varies from partial shade to full sun; has wet or saturated soil, standing water, or a high water table; generally populated with wetland and wet meadow species.



FERTILITY

Due to the potential for water contamination, lime and fertilizer are not recommended; however, when topsoil has been depleted or removed, we recommend the addition of organic matter (compost). Check soil pH and select species adapted to that pH.



SEEDING METHOD

Hand seed, broadcast seed, hydroseed, or drill seed when the water table is drawn down. It is not practical to seed any wetland where there is standing water or where severe flooding is likely to occur before germination. The same caution applies to mulching. Natural seed banks (seeds in wetland soils) often establish part of the vegetative cover.



Wetlands are massive nutrient filters that help clean our nation's water supply and protect rivers and oceans from pollution.

GROWING SEASON MAINTENANCE

FIRST GROWING SEASON

When feasible (the ground isn't too slippery or mucky to safely walk), post-planting maintenance will provide the best results for wet meadows and wetlands. Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a string trimmer. Trimming reduces competition by fast-growing weeds for sunlight and nutrients needed by slower growing perennial natives. Trimming should cease by mid-September.


Problem weeds should be hand pulled or spot sprayed with an approved aquatic herbicide, such as Rodeo® or Garlon® 3A.

SECOND & SUBSEQUENT GROWING SEASONS

Problem weeds, such as purple loosestrife, phragmites, Japanese knotweed, and reed canarygrass, should be hand pulled or spot sprayed with an approved aquatic herbicide, such as Rodeo® or Garlon® 3A.



Pickerelweed (*Pontederia cordata*) is a native wetland plant that provides food for aquatic wildlife.



ERNMX-131 OBL Wetland Mix.



WET MEADOW & WETLAND SITES SEED MIXES

ERNMX-120	OBL-FACW Perennial Food & Cover Wetland Mix
ERNMX-122	FACW Wetland Meadow Mix
ERNMX-128	Seasonally Flooded Wildlife Food Mix
ERNMX-131	OBL Wetland Mix
ERNMX-137	Specialized Wetland Mix for Shaded OBL-FACW Areas
ERNMX-138	Wildlife Food & Shelter Mix
ERNMX-175	Waterfowl Buffet Mix for Wetland Enhancement
ERNMX-176	Passive Acid Mine OBL Wetland Mix



Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.

PARTIALLY SHADED SITES

FAST FACTS

Partially Shaded Sites are characterized as having a relative lack of direct sunlight, typically caused by trees and shrubs that reduce their exposure. Examples: Woodland openings and sites associated with bioengineering installations.



SITE PREPARATION

These sites generally involve working around trees and shrubs while minimizing damage to trunks and roots. Undesirable vegetation must be controlled by tilling or direct spraying with glyphosate. Invasive weeds not addressed before establishment will be difficult to remove later. The soil needs to be loosened in order to establish seed-to-soil contact and dense leaf litter should be broken up with a rototiller. Seedlings can emerge from light leaf litter if planted at the proper depth. Light mulch or hydromulch can protect the seeds and soil until germination. Seeding and mulching around bioengineering material should take place immediately after installation. If installation cannot take place immediately after grading, temporary seeding and mulching are recommended.



HABITAT

Typically in moderate shade; many native species are adapted to moderate shade and the protective habitat around trees; shade tolerant native grass species, such as *Agrostis perennans* (Autumn Bentgrass), *Chasmanthium laxum* (Slender Woodoats), *Cinna arundinacea* (Wood Reedgrass), *Elymus hystrix* (Bottlebrush Grass), *Elymus riparius* (Riverbank Wildrye), *Elymus virginicus* (Virginia Wildrye), and *Panicum clandestinum* (Deertongue), provide early protection for emerging herbaceous species. Note: For understory of longleaf pine plantings, high biomass producing species, such as switchgrass, big bluestem, and indiagrass, should be avoided. Fire can be too hot for longleaf pine seedlings or trees when these species are burned.



FERTILITY

The addition of organic matter (compost) is most important. Check soil pH and select species adapted to that pH.



SEEDING METHOD

Hand seed, broadcast seed, or hydroseed. Use a garden rake, drag, or roll the surface to incorporate the seed into the soil 1/4"-1/2" deep. A seed drill may be used when sufficient room exists for operation.



Showy Ticktrefoil (*Desmodium canadense*) in a field.

GROWING SEASON MAINTENANCE

FIRST GROWING SEASON

Whenever canopy height (overall vegetation) reaches 18"-24", trim the meadow to 8" using a brush hog mower or string trimmer. Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

If bioengineering materials were used on the site, mowing should be above the new growth of these materials. Trimming should cease by mid-September.


Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Roundup®, Rodeo®, Garlon®, Garlon® 3A, Sonora™, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.

SECOND & SUBSEQUENT GROWING SEASONS

Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2"). This will allow the soil to warm more quickly, stimulating emergence and growth of herbaceous native plants and reducing invasion of woody undergrowth. In certain ecosystems, controlled burning by certified professionals can achieve the same results.

If bioengineering materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.

Problem weeds should be hand pulled or spot sprayed. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive, Canada thistle, and mugwort.



Woodland openings can be developed to provide crucial early successional habitat between old growth forests and adjoining grasslands.

SPECIAL CIRCUMSTANCES

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". Trimming should cease by mid-September.





Blunt Broom Sedge (*Carex scoparia*) provides food and cover for songbirds, ruffed grouse, and ducks.



Indianhemp (*Apocynum cannabinum*) is attractive to butterflies and songbirds.




Partridge Pea (*Chamaecrista fasciculata*) is found along wooded edges and provides great pollinator value in late summer and early fall.

PARTIALLY SHADED SITES SEED MIXES

ERNMX-132	Right-of-Way Non-Native Woods Mix
ERNMX-132-1	Right-of-Way Native Woods Mix with Annual Ryegrass
ERNMX-140	Partially Shaded Area Roadside Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.



STORMWATER MANAGEMENT SITES

FAST FACTS

Stormwater management facility sites are generally a best management practice integrated throughout land development projects which provide for a volume of water storage, infiltration, and evaporation mimicking the natural rate of runoff or groundwater recharge.



ABOUT STORMWATER MANAGEMENT SITES

Stormwater management sites (e.g., size, location, and depth) must be designed and constructed according to all applicable ordinances and under the direction of a qualified design professional. In addition to planting trees and shrubs, seeding with native plant species is an economical way to vegetate stormwater management sites. Erosion control fabric, mulch, or hydromulch is necessary to control erosion during and after construction.

Examples: Detention basins - temporarily impound run-off water, allowing for release at controlled rates; retention basins - stormwater management facilities with permanent impoundment or pool for improving water quality; bioretention basins and rain gardens - dynamic living microbiological systems that enhance retention, infiltration, and evaporation of run-off water while remaining attractive to wildlife.

PLANT SELECTION

In all stormwater management facilities, plants prevent erosion and slow water movement, hold or convert pollutants, enhance infiltration and evapotranspiration, and encourage wildlife. Plant species or mixes may be selected that meet critical objectives and extreme conditions of the site. Native grasses produce fibrous root systems that tolerate fast-moving water. Woody and herbaceous species add aesthetics, provide wildlife food and habitat, and assist with evapotranspiration while preventing erosion.

To avoid stand failure, select a seed mix appropriate for the site's hydrology (moisture status). If the site is chronically moist with long periods of inundation, a retention basin, FACW meadow, or OBL wetland mix is appropriate; if chronically moist with occasional inundation (immediately after a rainfall) and periodic drying out, a FACW meadow or riparian floodplain mix is appropriate; or, if typically dry except for a 12-72 hour period after a rainfall, a detention basin or rain garden mix is appropriate.



Stormwater basin utilizing ERNMX-126 Retention Basin Floor Mix - Low Maintenance.



Petrolia, Pennsylvania stormwater basin utilizing ERNMX-127 Retention Basin Wildlife Mix.

An urban rain garden catch basin utilizing ERNMX-180 Rain Garden Mix.

SITE PREPARATION

Prior to planting the site, invasive species, particularly those adapted to wet conditions, should be removed or sprayed using an approved herbicide by a licensed spray technician. Perennial weeds not addressed before establishment will be difficult to remove later. Normal vegetation can be worked into the topsoil which should be stockpiled until the final grade has been established.

With specifications and dimensions, on-site construction of the berm and outlets must be executed carefully to maintain structural integrity. The infiltration and plant growth areas should be loose and friable (easy to crumble), high in organic matter, and completed without compaction by heavy equipment. An excavator may be used to dig and drop each area of the bottom soil in a loose manner. Lime or compost can then be incorporated. The excavation machine does not move over the finished surface thereby avoiding unnecessary compaction. Native vegetation can be planted or seeded over this uneven absorbent surface.

SEEDING AND PLANTING METHODS

Seeding and planting should begin immediately upon completion of the structure while the soil is still friable and before weeds emerge. Plan seeding and planting before the basin is flooded or allow the basin to drain before seeding. Broadcast seed evenly over each unit by hand seeding or hydroseeding. Seeding rates are generally low (1/2 lb per 1,000 sq ft). If broadcasting seed, see Seeding Methods on p. 19. If the soil is dry, incorporate the seed into the soil with a garden rake. Oats, Japanese millet, or rye can provide temporary vegetation to protect the soil until permanent vegetation is established. Using such native species as *Elymus virginicus* (Virginia Wildrye) can create an intermediate cover that is succeeded by long-term native vegetation. Straw mulch or straw coconut mats are frequently used to control erosion and protect emerging seedlings from extreme temperatures and drying out. Mulch should be sparse to allow sunlight to reach the ground. If the site is a retention basin, refer to the Wet Meadow & Wetland Sites Establishment Guide, p. 42.

Transplanted seedlings and shrubs may need to be watered until they become well-rooted. Irrigating seeded areas is beneficial until seedlings become established.

GROWING SEASON MAINTENANCE

FIRST GROWING SEASON

Whenever canopy height (overall vegetation) reaches 18"-24", use a brush hog mower or string trimmer to trim the meadow to 8". Trimming reduces competition by fast-growing weeds for sunlight, water, and nutrients needed by slower growing perennial natives. A lawn mower is not recommended as the mower height will be too low and native seedlings will be killed.

If bioengineering or containerized woody materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo®, Garlon® 3A, or Milestone® (do not use Milestone® where standing water is present). Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after they have had two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

SECOND & SUBSEQUENT GROWING SEASONS

Prior to new spring growth reaching 2" (e.g., shortly after forsythia or redbud blooms), trim any material standing from the previous year close to the ground (approximately 2"). If the seed mix included sedges, trimming should be no lower than 2" above the crowns that produced the prior year's growth. This will allow the soil to warm more quickly, stimulating

emergence and growth of native plants and reducing the likelihood of shrub invasion.

If bioengineering or containerized woody materials were used on the site or seed of shrubs/trees were part of the mix, the site should not be trimmed after the establishment year.

Problem weeds should be hand pulled or spot sprayed with an approved herbicide, such as Rodeo®, Garlon® 3A, or Milestone®. Be vigilant in controlling vines or thorny plants if they were not part of the mix. These are more easily pulled early than after two to three months of growth. Examples include bindweed, blackberry, multiflora rose, mile-a-minute, and Japanese hops. Be equally vigilant in the control of other invasive species, such as autumn olive and Japanese knotweed.

SPECIAL CIRCUMSTANCES - SECOND GROWING SEASON

If there is a heavy infestation of ragweed or foxtail in the second growing season, trim the meadow to 8". If bioengineering or containerized woody materials were used, trimming should be above or around new growth of the plants. Trimming should cease after mid-September.

GENERAL MAINTENANCE

In addition to structural maintenance, siltation should be removed as needed. Close mowing throughout the growing season or extensive chemical use is not conducive to water quality improvement and wildlife habitat.

STORMWATER MANAGEMENT SITES SEED MIXES

ERNMX-126	Retention Basin Floor Mix - Low Maintenance
ERNMX-127	Retention Basin Wildlife Mix
ERNMX-128	Seasonally Flooded Wildlife Food Mix
ERNMX-154	Floodplain Mix
ERNMX-180	Rain Garden Mix
ERNMX-180-1	Rain Garden Grass Mix
ERNMX-180-2	Southeast Rain Garden Mix
ERNMX-183	Native Detention Area Mix


Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.



SOUTHEASTERN U.S. SITES

FAST FACTS

Southeastern U.S. sites include regions within Alabama, the Carolinas, southern Virginia, Georgia, Mississippi, and Florida. Sites in the Southeast typically have sandy or clay-rich soils subject to drought. These characteristics necessitate careful attention to timing and preparation. Examples: Coastal Plain soils, Piedmont, and sandy soils (refer to Upland & Meadow Sites Establishment Guide, p. 23).



A Florida meadow.

SITE PREPARATION

If the site was previously a lawn or crop field to which herbicides were applied, it is important to allow the appropriate time interval for the herbicide residues to break down prior to planting. Some herbicide residues can prevent seedlings from germinating.

Competition from invasive or undesirable vegetation is the most limiting factor in upland meadow preparation. Prior to planting, all such vegetation must be fully controlled. Typical control strategies include repeated tilling, smothering with black plastic, or herbicides. While any of these methods may control existing weeds, they will not kill all weed seeds lying dormant in the soil. Seeds of such species as velvetleaf and pokeweed may germinate many decades after the species last flowered on the site.

When using the tillage strategy, a site is disc harrowed every two weeks for one to two months. The underlying principle of this process is that the root system of perennial species will be worn out to the point of killing the species. In addition, tillage will stimulate germination of some dormant weed seeds that will be killed with subsequent tillage. Planting should not take place until perennial species are completely killed.

Black plastic may also be used to kill weeds. It may be laid across tilled or untilled soil and anchored down by burying the edges in soil or laying boards or bricks across the surface. This strategy should be utilized during a growing season when the intent is to fall plant the same year or spring plant the following year.

Use of an approved herbicide, such as glyphosate (Roundup® or Rodeo®), by a licensed spray technician is the most common and least time-intensive method for the control of existing vegetation. Since herbicides are most effective on actively growing plant tissues, they are very effective on new growth in the spring. Spraying should begin when growth reaches 6". A follow-up spray application one to two weeks later will address skips or persistent species. If substantial plant tissue remains on the surface following a full kill by herbicides, a close mowing, tillage, or burning may be necessary to achieve good seed-to-soil contact.

To prevent reinfestation of some weed species, use of an appropriate selective herbicide in conjunction with a seed mix tolerant of that herbicide may be necessary.

SANDY SOILS

Sandy soils behave differently under cultivation than those containing clay. It is essential to plant seed 1/2" deep into a firm seedbed with a seed drill if possible (Eastern Gamagrass should be planted 1" deep). Truax and other similar drills can accommodate a variety of seeds and have been proven effective in the Southeast. High sand content in these soils makes broadcasting seed less effective due to poor seed-to-soil contact. Seedbeds should be firmed to where one does not sink past the sole of his/her shoe when walking the prepared site. Soil amendments may be added as necessary to maintain proper levels of organic matter and achieve a pH of at least 5.5.

CLAY-RICH SOILS

Without topsoil, soils containing high clay levels can be as hard as brick and pose a formidable challenge for successful cultivation. These soils are extremely low in organic matter which allows the small clay particles to settle and become compacted after a rain event. They are often iron-rich, leading to a distinctive red color. To prevent the clay from hardening after a rain from which seedlings cannot emerge, increase soil organic matter by incorporating 1"-2" of well-decomposed organic matter or compost and working it into the top-most soil prior to planting using a tiller, harrow, disc, or similar implement. Cultivating the top 6"-8" of soil will aid in root development of emergent seedlings and allow some percolation of rainwater that would otherwise run off the surface with little to no infiltration and carry the seed away with it. These initial preparations are critical for the successful establishment of native plants in this challenging soil. Since soil compaction is minimized, drilling seed 1/4"-1/2" deep is the preferred planting method. Even with additional organic matter, this clay-rich soil will compact easily; therefore, operating heavy equipment over the planted site should be avoided.



HABITAT

Southeastern sites have a longer growing season; therefore, plants native or adapted to the region should be selected; planting from November to March is ideal (when possible) as temperatures are adequate and rain events are frequent; if irrigation is available, planting can continue into the later months of spring and early summer.



FERTILITY

With the exception of organic matter, natural fertility is generally adequate. Check soil pH and, if necessary, add lime to achieve a pH of at least 5.5.



SEEDING METHOD

Drill seeding is recommended; however, broadcast seeding is an alternative preceded by rolling or tracking.



Spiked Wild Indigo (*Baptisia albenscens*) in a South Carolina pine flatwoods.



Quail-friendly planting featuring Partridge Pea (*Chamaecrista fasciculata*), Beaked Panicgrass (*Panicum anceps*), Blackeyed Susan (*Rudbeckia hirta*), and Little Bluestem (*Schizachyrium scoparium*).

GROWING SEASON MAINTENANCE

MAINTENANCE

Refer to Upland & Meadow Sites maintenance, p. 23)

When spot spraying in soils with low organic matter and high sand levels, begin with lower than recommended concentrations of herbicides for weed control to avoid valuable crop burnout. Chemical breakdown of many herbicides is achieved via soil microbes that generally feed off organic material. With less organic material available in the soil, there will be a smaller population of microbes that may result in longer periods of exposure to the active ingredients in herbicides.

SOUTHEASTERN U.S. SITES SEED MIXES

ERNMX-169	Southeast Annual & Perennial Wildflower Mix
ERNMX-187	Southeastern U.S. Roadside Native Mix

Mix formulations are subject to change without notice depending on the availability of existing and new products. While the formula may change, the guiding philosophy and function of the mix will not. See "Disclaimer," p. 15. For "Expectations of Native Species," see p. 12.



These mixes are used in well-drained soils with full sun and provide food and/or cover for wildlife. Meadow and wildflower mixes provide food for insects, including native pollinators.



OBSTACLES TO A SUCCESSFUL ESTABLISHMENT

FAST FACTS

Over the past 20 years, we have had the opportunity to visit sites or see photographic evidence of many native meadows that failed to establish successfully. As a result, we have found some repetitive themes relating to these failures - site conditions, weeds, excessive cover crop seeding rates, or use of an inappropriate cover or companion crop.



Stand failure in central Pennsylvania caused by excess application of Annual Ryegrass (*Lolium multiflorum*).

OBSTACLES TO A SUCCESSFUL ESTABLISHMENT

POOR SITE CONDITIONS

- Poor pre-plant weed control: Native species require bare ground to germinate and grow and will not establish well in a site already vegetated with weeds or lawn.
- Presence of excessive organic matter used as mulch: Mulch prevents good seed-to-soil contact. A seed may germinate but its radicle (first root) may be unable to find moisture and the seedling dies. Microorganisms that decompose mulch or compost consume nitrogen, a nutrient essential to plant growth. This results in a nitrogen-deficient environment in which a seedling will grow poorly or not at all. If using compost, be certain it has decomposed to where the parent material is unrecognizable.
- Soil compaction: If the tip of a crowbar or piece of rebar cannot be pressed 3" deep into the soil, there is compaction which will result in a very poor or non-existent stand. For highly compacted soils, consider broadcasting 1"-2" of well-decomposed compost across the site followed by rototilling the compost in 6" deep.
- Herbicide residues, including pre-emergent herbicides, from the previous year: Generally, occurs on sites that were previously lawns with a lawn service contract for weed control within 12 months of planting. Also problematic is when the site was a crop field to which herbicides, such as Atrazine (atrazine), Princep® (simazine), Milestone® (aminopyralid), Sonora™ (clopyralid), Resolve® Q (rimsulfuron), Cimarron® (metsulfuron methyl), or Synchrony® XP (chlorimuron ethyl), were applied within 12 months of planting.



Stand failure caused by soil compaction at a landfill in northwestern Pennsylvania.

- Excessively high or low soil pH: For many plant species native to eastern North America, availability of many nutrients essential to plant growth is limited in soils with pH less than 4.5 or greater than 7.5. Ideal pH is 5.5-7.0. When pH is outside this range, species tolerant of the site's pH should be chosen (pH can be raised with lime or lowered with sulfur).



Canada thistle in a wetland restoration site.

- Drought within two to six weeks after planting: Seeds cannot germinate without water and seedlings do not grow/survive without water. In drought-prone areas, we recommend planting between late October or when soil temperatures are less than 55°F (13°C) at 3" deep and in spring when forsythia or redbud bloom.
- Erosion (on steep slopes): Failure to use erosion control blankets or toe them in at the top of the slope. Erosion is also a problem on slopes where final tracking of the soil was not perpendicular to the slope, resulting in the seed being washed down to the bottom of the slope.
- Presence of crownvetch, sericea lespedeza, trefoil, tall fescue, bahiagrass, bermudagrass, white Dutch clover, alsike clover, bindweed, mile-a-minute, Japanese hops, kudzu, or invasive shrub species not controlled prior to planting or volunteered from dormant seed when the soil was prepared for planting. These species can smother desirable but slower growing perennial natives.
- Use of borrowed topsoil infested with seeds of invasive species.

- Wildlife: Geese can eat seeds, seedlings, and mature plants while deer can be devastating to some wildflowers. If up to five deer are regularly observed in the area, it may be wise to plant a deer-resistant mix. If the deer population is sufficient to make growing a garden or fruit trees difficult, it may be impossible to grow wildflowers on the site. The same is true if a deer population of 20 or more is regularly observed in the area.
- Lack of proper maintenance: Letting annual ryegrass cover crop or weeds, such as foxtail or ragweed, smother native seedlings during the first full growing season. Avoid this by trimming the meadow to 8" whenever growth exceeds 18"-24".
- Failure to control invasive or problem species prior to planting or after germination: If there are vines or spiny plants in the landscape that were not planted, be vigilant in controlling them.

APPLICATION OF A COVER CROP AT AN EXCESSIVE RATE

Issues with an excessive rate of cover crop have generally been confined to the use of annual ryegrass. Excessive annual ryegrass applied to sites planted with native species smothers growth of the native meadow. We have not had complaints when annual ryegrass was used at 10-12 lb per acre.

USE OF AN INAPPROPRIATE COVER OR COMPANION CROP

We do not recommend the use of bahiagrass, bermudagrass, or tall fescue as cover or companion crops in our native meadow mixes. Bahiagrass or bermudagrass as a cover crop will be impossible to control prior to new growth of perennial native species and will likely smother out the native species when used as a companion crop.

We do not recommend the addition of the following legumes to native meadows as they also tend to take over and smother out native species: *Lespedeza cuneata* (Sericea Lespedeza), *Coronilla varia* (Crownvetch), *Lotus corniculatus* (Bird's Foot Trefoil), *Trifolium pratense* (Red Clover), or *Trifolium repens* (Ladino Clover). If already present on the site, control these species by spot spraying with Roundup® (glyphosate).

If a legume is desired in a meadow mix, we recommend such native legumes as *Baptisia alba* (White Wild Indigo), *Baptisia albescens* (Spiked Wild Indigo), *Baptisia australis* (Blue False Indigo), *Baptisia tinctoria* (Yellow False Indigo), *Chamaecrista fasciculata* (Partridge Pea), *Chamaecrista nictitans* (Sensitive Pea), *Desmodium canadense* (Showy Ticktrefoil), *Desmodium paniculatum* (Panicked Ticktrefoil), *Lespedeza capitata* (Roundhead Lespedeza), *Lespedeza frutescens* (Shrubby Bushclover), *Lespedeza virginica* (Slender Bushclover), *Senna hebecarpa* (Wild Senna), and *Senna marilandica* (Maryland Senna).



An abundance of Canada geese can decimate seeds and seedlings.



Green foxtail smothered native seedlings during the first full growing season of this planting.



Japanese hops can quickly overtake a native planting if not controlled early.

WHAT IS THE APPROPRIATE COVER CROP FOR A MEADOW?

We recommend the following cover crops and seeding rates:

- **Grain Oats:** 30 lb per acre; planted January through July (areas north of the Mason-Dixon Line), January through April (areas south of the Mason-Dixon Line).
- **Grain Rye:** 30 lb per acre; planted August through December (areas north of the Mason-Dixon Line), September through December (areas south of the Mason-Dixon Line), and September through April (moist sites).
- **Annual Rye:** 10-12 lb per acre; planted year-round (dry sites).
- **Brown Top Millet:** 10 lb per acre; planted May through August (dry sites south of the Mason-Dixon Line).
- **Japanese Millet:** 10 lb per acre; planted May through August (wet sites).

These seeding rates are based on our experience with native meadows as well as our desire

to establish strong, individual native plants. Planting cover crops that are too aggressive or thick diminishes the long-term viability of the perennial meadow plants. We have concluded that annual small grains, such as oats and rye, are the best cover crops or companion crops to plant with native seedlings when there is a need. Grain cover crops can reduce competition from aggressive weeds because they grow quickly and reduce the potential for erosion by providing quick cover. We generally do not recommend annual ryegrass as it is too aggressive and volunteer seedlings can be persistent. When using annual ryegrass with native species, do not exceed 12 lb per acre.

PROBLEM WEEDS FOR UPLAND MEADOWS

In much of our market area, crabgrass, giant foxtail, green foxtail, and ragweed can smother a meadow in the establishment year. If overtaken by these weeds, use a brush hog mower or string trimmer to trim the meadow to 8". Trimming below 4" will kill seedlings of many native species. A lawn mower is not recommended.



Ernst Seeds

BIOMASS OVERVIEW



BIOMASS OVERVIEW

Ernst Seeds is one of the largest switchgrass seed producers in the country, having more than 30 years' experience in the establishment, management, and harvest of native warm season grass seed and biomass.

Switchgrass, as well as other native warm season grasses, has attracted much attention as a potential source of alternative energy and sustainable fiber due to the following:

- Native warm season grasses are perennial.
- Native warm season grasses thrive in marginal soil conditions too wet or dry for traditional crops.
- Native warm season grasses require minimal nutrient input.
- Native warm season grasses are efficient in converting sunlight to usable biomass.
- Native warm season grasses have proven soil, water, air, and wildlife benefits.

Biomass production from switchgrass can vary greatly from one region to another. It is important to select switchgrass varieties well-suited to the growing conditions of your area. Please contact us and we will be happy to make recommendations.

A mix of switchgrass varieties adapted to your area can better acclimate to seasonal variation and soil conditions than a single variety. Diverse genetic material will allow the overall stand to thrive in a wider range of moistures, soil types, disease pressures, and weather.

Commonly, mixes that include other native grasses, such as *Andropogon gerardii* (Big Bluestem), *Sorghastrum nutans* (Indiangrass), *Schizachyrium scoparium* (Little Bluestem), *Panicum amarum* (Coastal Panicgrass), and *Spartina spp.* (Cordgrass spp.), may create a biomass product that will satisfy many conservation program requirements while also being able to be marketed.



Switchgrass makes a highly effective livestock forage and is increasingly used as a stand-alone grazing stock and in diverse native grazing seed mixes.

SELECTING THE RIGHT BIOMASS VARIETIES

Our supply of switchgrass seed comes from various sources, including our licensing of the varieties produced by intensive breeding programs at numerous institutions and regional populations made available from USDA plant materials centers. The regional populations have minimal genetic improvement for general physical characteristics and have been adequate

for decades for erosion control, wildlife plantings, and in the Conservation Reserve Program (CRP). The new varieties, including 'Mt. Airy', 'BoMaster', 'Timber', 'Liberty', 'Independence', 'Shawnee', and RC Chippewa, have significant yield improvements and were bred with a focus on forage and biomass production.



FORAGE

As with several other native warm season grasses, switchgrass can produce high-quality forage. Used in a system of rotational grazing, switchgrass allows for robust growth during hot summer months. University of Tennessee findings suggest that the nutrient content of this forage can be as high as 16%-17% crude protein.

Ground switchgrass straw is experiencing increased use as a forage extender in livestock feeds in that it works to increase bulk and dilute protein in operations with sources of high-protein feed.

NUTRIENT RUN-OFF PREVENTION

Switchgrass has extensive roots, growing as deep as 5'-6'. In addition to serving as a superior soil stabilizer in erosion control, switchgrass and its root system form a tremendous ecological filter, soaking up such nutrients as nitrogen and sequestering carbon dioxide. Use of switchgrass as a buffer or part of a riparian system between agricultural activity and watersheds is seen by many as one of the best methods for protecting these priceless resources.



Calvin Ernst with a mature stand of Switchgrass (*Panicum virgatum*).




As a riparian buffer, the extensive root system and nutrient-filtering qualities of switchgrass make it a powerful option.

POULTRY AND DAIRY BEDDING

Numerous studies have shown ground switchgrass to be easy on the pads of chicken feet, highly absorbent, and may represent a benefit over other beddings in the reduction of ammonia. From a cost perspective, producers can grow switchgrass on their own marginal ground, then harvest and process it for their own bedding uses. In addition to helping to control noise and water pollution, switchgrass can aid in making areas of marginal ground productive by supplying sustainable bedding.




Chick nestled in ground Switchgrass bedding.



Ground switchgrass is gaining popularity as an effective, readily accessible, and inexpensive livestock and poultry bedding material.

For information on preparing a field for the establishment and production of native biomass, please visit the following pages. For more information, please consult the Native Biomass FAQs at www.ernstseed.com.

BIOMASS PRODUCTION SITES



FAST FACTS

Biomass Production Sites are sites planted with warm season grasses specifically for harvesting or grazing livestock on the biomass (vegetative matter). Special focus is given to the fact that these perennial crops may be harvested from the site for a decade or more without replanting.

ADAPTING A FIELD FOR BIOMASS PRODUCTION

Every field has unique characteristics to be considered when establishing perennial native biomass species. These include soil type, hydrology, pH, fertility, erosion/run-off potential, compaction, existing vegetative cover, previous crop history, and harvest methods.

While natural soil type cannot be changed, native warm season species can tolerate virtually any soil type. Switchgrass can survive in a wide range of soil moisture. As is the case with row crops or alfalfa, biomass productivity will be directly related to soil quality.

Switchgrass can tolerate soil pH of 5.0-8.0 but will produce well at 6.0. Soil pH below 6.0 should be corrected with the addition of lime according to soil test recommendations.

Soil fertility is a function of the available nutrients that can be used by the plant. Warm season grasses (switchgrass in particular) can be more productive at lower fertility levels than row crops or alfalfa. Soil tests are required to determine soil fertility levels. Fertility levels referred to as moderate are generally adequate for biomass production. Fertilizer is not recommended for soils with moderate fertility levels. Adding nitrogen in the second and subsequent years is recommended based on expected yields.



Switchgrass grows through corn stubble, reducing its need for weed control.

Fields having a history of good weed control are the easiest to convert to native warm season grasses, such as those planted in corn or soybeans. Fields in conventional hay or pasture are somewhat more difficult to seed and require Roundup® to kill the cool season grasses as well as minimum tillage to work thatch into the subsoil.

No till or minimum till are the most effective means of seeding new warm season grasses. A limiting factor of no till is surface crop residue that prevents proper seed-to-soil contact, shades the germinating seedlings, and/or creates a nitrogen deficiency during decomposition. Each situation requires customized tillage and herbicide considerations.

Tilling a field going to warm season grasses corrects surface roughness and incorporates

crop residue before planting. The field surface must be smooth enough to spray, plant, mow, and bale.

Fields with perennial or invasive weeds that have not been mowed during previous growing seasons are the most difficult to prepare for native grass establishment.

Perennial vegetation of grasses and broadleaf weeds must be controlled prior to planting. Mowing or burning existing vegetation will produce new vegetative growth. Roundup® and a systemic broadleaf herbicide can then be sprayed to effectively kill undesirable species. Identify weeds present and use label rates to control weeds with one or more applications. Once controlled, seeding like a conventional hayfield may proceed.



Planting switchgrass in rows makes it easier to identify the plant in emerging stands.



A prescribed burn of existing vegetation prior to establishment of warm season grasses.



Application of Roundup® and systemic broadleaf herbicides is an important step in controlling weeds.



BIOENGINEERING MATERIALS

BIOENGINEERING MATERIALS

SOIL BIOENGINEERING

Soil bioengineering is the term for using plant material to arrest and prevent slope and streambank failure and erosion. The roots and stems serve as structural and mechanical elements in a slope protection system. Live cuttings and rooted plants are embedded in the ground in various arrays to serve as soil reinforcements, hydraulic drains, and barriers to earth movement. Once established, this living material effectively controls several stabilization and erosion control problems by binding the soil with its root system and creating a natural vegetative cover. Bioengineered sites are self-repairing and have the advantage of blending with natural surroundings.



Live stakes installed and sprouting new growth.

BIOENGINEERING MATERIALS & STREAM RESTORATION TECHNIQUES

Ernst Seeds is an experienced producer of common and specialized live soil bioengineering materials. We understand the unique needs of bioengineering site construction. Our material is grown, processed, and delivered to minimize on-site installation labor and maximize survival and quick establishment.

HELPFUL TIPS FOR A BIOENGINEERING PROJECT

Ernst Conservation Seeds' bioengineering products are dormant live material. Therefore, if installation cannot take place immediately upon arrival at the site, these products must be stored properly. Place in a cool, wet place out of direct sunlight, such as under straw or burlap. Open any pallets, boxes, and plastic bags so the

material can be watered thoroughly. Do not allow the material to dry out. Soaking before planting significantly increases survival and growth rate.

For best survivability, the material should be planted during the dormant season, November 1st-April 30th. We do not guarantee any of our bioengineering material from May 1st-October 31st.

Overseeding and mulching a completed bioengineered project with the appropriate seed mixes protects the soil surface from erosion while adding biodiversity to the site.

EXCELLENT MIXES FOR THIS PURPOSE

ERNMX-137	Specialized Wetland Mix for Shaded OBL-FACW Areas
ERNMX-138	Wildlife Food & Shelter Mix
ERNMX-178	Riparian Buffer Mix



PLACING AN ORDER


Please call for a quote as each bioengineering job and material selection is unique to a project.



TURNAROUND TIME

Since we custom cut your order, we require at least two-weeks' notice when bioengineering orders are placed.

STREAMBANK DIAGRAM



WE PRODUCE BIOENGINEERING MATERIALS IN THESE FAST-ROOTING SPECIES:

BOTANICAL NAME	COMMON NAME	PLANT TYPE
<i>Cephalanthus occidentalis</i>	Buttonbush	Native Shrub
<i>Cornus amomum</i>	Silky Dogwood	Native Shrub
<i>Cornus sericea</i>	Red Osier Dogwood	Native Shrub
<i>Salix amygdaloides</i>	Peachleaf Willow	Native Tree
<i>Salix discolor</i>	Pussy Willow	Native Tree
<i>Salix exigua ssp. interior</i>	Sandbar Willow	Native Shrub
<i>Salix lucida</i>	Shining Willow	Native Shrub
<i>Salix nigra</i>	Black Willow	Native Tree
<i>Salix purpurea</i>	Streamco Willow	Naturalized Shrub
<i>Salix sericea</i>	Silky Willow	Native Shrub
<i>Salix x cottetii</i>	Bankers' Dwarf Willow	Naturalized Shrub
<i>Sambucus canadensis</i>	Elderberry	Native Shrub
<i>Viburnum dentatum</i>	Arrowwood	Native Shrub
<i>Viburnum lentago</i>	Nannyberry	Native Shrub

For more information on the species listed above, refer to *Partially Shaded Sites*, p. 46.

Live Stake and Branch Layering Cross-Sections courtesy of United States Department of Agriculture, Natural Resources Conservation Service (NRCS), *Engineering Field Handbook*, December 1996, Chapter 16, "Streambank and Shoreline Protection", pp. 16-13 and 16-20.

Special thanks to John McCullah, *Salix Applied Earthcare*, for allowing us to use the information in his *Bio-Draw* software. More information is available at www.biodraw.com.

LIVE STAKES


Stakes are dormant, live woody cuttings of a species with the branches trimmed off. Live staking performs an important function in creating a root mat that stabilizes the soil by reinforcing and binding soil particles together. Stake establishment also improves aesthetics and provides a habitat for wildlife. Live stakes may be used on their own to secure other bioengineering materials or as an anchor for erosion control and geo-fabric. Stakes or poles may also be inserted or driven through openings in rock structures, such as gabions, riprap, and other retaining structures.

INSTALLATION NOTES: Install stakes during their dormancy (late fall to early spring). Do not allow the material to dry out. Soaking before planting significantly increases survival and growth rate. Drive a pilot hole into firm soil and plant at right angles (buds oriented up) with at least two-thirds of its length underground. Plant stakes randomly or 3'-6' apart on triangular spacing. Tamp the soil down around the cuttings before watering. Irrigation may be necessary if a long dry spell or hot weather is expected following installation.

SIZES: 1/4"-1" diameter; 2'-4' lengths.

- A. Geotextile fabric
- B. Toe protection
- C. Live cutting 1/2" to 1 1/2" dia.
- D. 2' to 3'
- E. 2' to 3' (triangular spacing)
- F. Dead stout stake
- G. Erosion control fabric
- H. Stream bank

CROSS-SECTION DIAGRAM (not to scale)




BRUSH (BRANCH) LAYERS

Brush (branch) layers are living branches placed on a terrace along the contours of a streambank and interspersed between layers of soil. This technique is used to repair a slump or gully and is most effective for revegetating scour holes.

INSTALLATION NOTES: Brush (branch) layers are placed on terraced benches with two-thirds of the basal material tilted into the slope and covered with soil. Branches should protrude beyond the face of the terrace. Before installing, soil terraces can be additionally protected by putting down geotextile fabric. Starting at the bottom of the slope, secondary brush (branch) layers may be added every 3'-4' proceeding up the slope. Straw mulching the finished surface is recommended for moisture retention and additional erosion control. Planting should be during the dormant season.

SIZES: 3 linear ft per bundle, 3'-6' lengths, 28-36 branches per bundle.





BRUSH MATTRESSES

Brush mattresses are living branches layered 1-2 branches thick in a crisscross pattern on a streambank to form a living ground cover. The mattress formed protects the bank surface until the branches root and native vegetation is established. This living system normally roots in the entire bank face, encouraging natural infiltration and immediately acting as a sediment trap.

INSTALLATION NOTES: Place material with the basal ends located toward the bottom of the slope. Using 3' square spacing, drive dead wedge stakes into the plant material. Stretch wire or biodegradable twine or rope diagonally around the dead stakes and finish driving the stakes in to tighten the wire, twine, or rope and secure the mattress. Place a thin layer of soil over the entire area to encourage rooting. Irrigation is necessary immediately following installation.

SIZES: 1/4"-3" diameter x random length x 1-2 or 3-4 branches thick.





A 5' Wattle, ready to ship.

WATTLES/FASCINES

Wattles or fascines are living branches bound together in long tubular bundles. When placed in shallow trenches across the slope of a bank, these structures provide protection from erosion and create a sediment trap that provides immediate bank support even prior to root growth. Once established, this live rooting material grows into a living fence-like erosion barrier. Within one growing season, roots and shoots grow along the entire length of the structure and quickly stabilize the bank.

INSTALLATION NOTES: This technique is simple, effective, and installed with little site disturbance. Material is placed in 6" wide trenches on banks or slopes parallel to the stream contour and partially covered with soil. Wedge-like dead stakes secure them in place at 2'-3' intervals. Live stakes may also be used in conjunction with dead stakes to secure the material. Straw mulching the site following installation retains moisture and reduces surface erosion. Irrigation is necessary after installation if the soil is dry.

SIZES: 5' or 6' lengths are recommended for ease of handling; available in the following diameters: 4"-5", 6"-8", 9"-12" (custom lengths and diameters are available).



Dead Wedge Stakes

DEAD WEDGE STAKES

Dead wedge stakes are pieces of hardwood cut into long wedges to secure wattles, brush mattresses, and other applications of soil bioengineering and erosion control measures.

SIZES: 1-1/4" x 3-1/4" x 2-1/2' long.

LIVE WHIPS

Whips are slender, live woody shrub material well-suited for very moist areas of stream edges, commonly used in conjunction with gabion structures, riprap, and geo-fabric.

INSTALLATION NOTES: Push whips into the ground as far as they will go without breaking. At least two-thirds of the whip should be covered with soil. Whips may be installed either by laying them on an angle or planting them erect in the soil. When using whips with hard structures, be sure they are long enough to reach into the soil and moisture behind or below the structure. (Example: If installing whips through riprap, consider the 3' depth; therefore, install a 6' whip at least 2' into the moist soil behind the stone and 1' above the surface of the riprap).

SIZES: 3/8"-1" diameter; 4'-6' lengths.



ERNST SEEDS SPECIFIC

FAST FACTS

We will identify, collect, and propagate new species and ecotypes that meet our clients' needs from eastern Canada to the southeastern United States. Our native seeds are produced from species considered to be the most significant foundation of an effective native restoration or reclamation project. To ensure that our customers receive a quality product, all our seeds are harvested, conditioned, and tested under the highest quality standards.

In addition to 10,000 acres of native seed production at Ernst, we also supply seed from some of North America's top producers and collectors.

Ernst Seeds and its suppliers collect, grow, and process all our products in an ecologically sustainable and renewable manner.



Acorus americanus
SWEETFLAG

Native
Rhizomatous species; sometimes misnamed *Acorus calamus*, an introduced species; provides food and cover for wildlife.

HABITAT: Swamps, shallow pond water, wet meadows.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
pH: 5.6-7.2
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 70,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Agrostis perennans
AUTUMN BENTGRASS

Native
Rhizomatous species; adapted to mesic to dry areas, including steep slopes and forests with dappled to full sun; provides winter grazing for wildlife.

HABITAT: Dry open ground, areas in light shade.

CHARACTERISTICS:
Height: Up to 3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5.5-7.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 8,000,000
Seeding Rate: Up to 30 lb per acre alone; up to 2.4 lb per acre in a steep slope mix

Herbaceous Perennial



Agrostis alba
REDTOP

Naturalized
Rhizomatous sod-forming species; produces quick cover on road banks and diversion ditches for erosion control; adapts well to pipeline restoration; provides food for wildlife.

HABITAT: Coastal marshes, roadsides, open ground; establishes well in moist soils.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4.5-8
Bloom Period: Summer

Approx Seeds Per Lb: 4,851,000
Seeding Rate: Up to 25% of a mix; up to 40 lb per acre alone

Herbaceous Perennial



Agrostis stolonifera
CREEPING BENTGRASS

Naturalized
Stoloniferous sod-forming grass; used on reclamation sites, lawns, and golf course putting greens; also used for soil erosion control.

HABITAT: Wet meadows, shores, wet to dry fields, roadsides.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.1-7.5
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 6,130,000
Seeding Rate: Up to 5% of a mix; of a fine fescue mix; up to 40 lb per acre alone

Herbaceous Perennial



Agrostis hyemalis
WINTER BENTGRASS


Native
Bunchgrass; good for partially drained soils on moderately shaded roadsides; provides winter grazing for wildlife.

HABITAT: Roadsides, meadows, fields, moist to dry open, sterile soils.

CHARACTERISTICS:
Height: Up to 2.5 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5-7.5
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 8,500,000
Seeding Rate: Up to 30 lb per acre alone; up to 2.4 lb per acre in a steep slope mix

Herbaceous Perennial



Agrostis tenuis
COLONIAL BENTGRASS

Naturalized
Rhizomatous species; provides erosion control; naturalized in many areas of the northeastern U.S. where sheep were grazed.

HABITAT: Cultivated in pastures, lawns, dry open ground, along roadsides.

CHARACTERISTICS:
Height: Up to 2.2 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4.9-7.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 6,130,000
Seeding Rate: Up to 5% of a mix; of a fescue mix; up to 40 lb per acre alone

Herbaceous Perennial



Alisma subcordatum
MUD PLANTAIN

Native

Grows quickly in early spring; produces seed in the fall; ideal for vernal pools; provides food for pheasants and waterfowl.

HABITAT: Marshes, stream sides, muddy shores, pond margins, shallow water.

CHARACTERISTICS:

Height: Up to 3.3 ft.

Minimum Root Depth: 8 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

Pollinator Value: Low

pH: 5-7

Bloom Period: Spring/Fall

Flower Color(s): White

Approx Seeds Per Lb: 825,000

Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Andropogon gerardii
BIG BLUESTEM

Native

Warm season bunchgrass; used for erosion control in sand and gravel pits, mine spoil, and on roadsides; contributes to diversified biomass production; high quality livestock forage; provides food and cover for wildlife.

HABITAT: Riverbanks, roadsides, meadows, open woods, savannas, tallgrass prairies.

CHARACTERISTICS:

Height: Up to 8.1 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 6-7.5

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 144,000

Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Allium cernuum
NODDING ONION

Native

Bulb-forming species; ideal ornamental for rock gardens; readily reseeds itself.

HABITAT: Dry rocky slopes, wood borders, rocky banks, prairies; often limestone outcrops.

CHARACTERISTICS:

Height: Up to 2 ft.

Shade Tolerance: Moderate

Drought Tolerance: Unknown

Pollinator Value: High

Bloom Period: Summer/Fall

Flower Color(s): White, Pink

Approx Seeds Per Lb: 123,000

Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Andropogon ternarius
SPLITBEARD BLUESTEM

Native

Warm season bunchgrass; white seedheads stand out in upland habitats of the southern half of the U.S.

HABITAT: Pinelands, sandhills, old fields.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Shade

Drought Tolerance: High

pH: 4-7.5

Bloom Period: Spring/Summer

Approx Seeds Per Lb: 216,000

Seeding Rate: Up to 10% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Alopecurus arundinaceus
CREeping FOXTAIL

Naturalized

Rhizomatous species; ideal for pipeline restoration where wildlife is desired; provides seed, forage, and cover for wildlife and domestic animals.

HABITAT: Wet hay meadows, margins of lakes and ponds, waterways.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

pH: 5.5-8.4

Bloom Period: Spring

Approx Seeds Per Lb: 786,000

Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Andropogon virginicus
BROOMSEEDGE

Native

Warm season bunchgrass; historically used to make brooms; stiff straw stays erect on road cuts during the winter; provides cover for wildlife.

HABITAT: Pastures, open woods, old fields, hillsides, open ground with dry infertile soils.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 4.9-7

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 800,000

Seeding Rate: Up to 2.5% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Anthoxanthum odoratum
SWEET VERNALGRASS

Naturalized

Dense, clump-forming cool season bunchgrass; sweet fragrance of freshly mowed hay when crushed.

HABITAT: Open fields, meadows, roadsides; grows primarily in poor soils.

CHARACTERISTICS:

Height: Up to 2.4 ft.
Shade Tolerance: Full Sun
Drought Tolerance: High
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 738,000
Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Bromus inermis
SMOOTH BROME

Naturalized

Sod-forming cool season grass; provides food for livestock.

HABITAT: Roadsides; grows best in well-drained fine-textured soils.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.5-8
Bloom Period: Summer

Approx Seeds Per Lb: 143,000
Seeding Rate: Up to 15% of a mix

Herbaceous Perennial



Bouteloua curtipendula
SIDEOATS GRAMA

Native

Fast-emerging warm season bunchgrass for upland meadows where sight lines are important; used for surface mine revegetation, erosion control, and as a warm season companion crop; high winter forage value for wildlife and livestock.

HABITAT: Dry woods, dry calcareous clearings, dry prairies, sandhills.

CHARACTERISTICS:

Height: Up to 3.2 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.5-8.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 159,000
Seeding Rate: Up to 35% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Calamagrostis canadensis
CANADA BLUEJOINT

Native

Rhizomatous grass; provides food and cover for deer, muskrats, and moose.

HABITAT: Swamps, wet meadows.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4.5-8
Bloom Period: Summer

Approx Seeds Per Lb: 3,837,000
Seeding Rate: Up to 0.5% of a mix; up to 2 PLS lb per acre alone

Herbaceous Perennial



Bouteloua gracilis
BLUE GRAMA

Native

Fast-emerging warm season bunchgrass; used in dry highway medians, recreation area plantings, and in pure stands for erosion control; provides food for wildlife.

HABITAT: Dry prairies, sandhills.

CHARACTERISTICS:

Height: Up to 2 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 6.6-8.4
Bloom Period: Summer

Approx Seeds Per Lb: 724,000
Seeding Rate: Up to 10% of a mix; up to 6 PLS lb per acre alone

Herbaceous Perennial



Carex albolutescens
GREENWHITE SEDGE

Native

Tufted sedge; provides food for wildlife.

HABITAT: Low fields, meadows, marshes, floodplain forests, thickets.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.3-7.8
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,000,000
Seeding Rate: Up to 20% of a mix

Herbaceous Perennial



Carex comosa
COSMOS SEDGE

Native
Bunch-type sedge; provides food and cover for wildlife.

HABITAT: Swamps, marshes, swales.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.6-7.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 480,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex grayi
GRAY'S SEDGE

Native
Bunch-type sedge; may be used for ornamental purposes in shaded areas; provides food and cover for wildlife.

HABITAT: Swamps, wet woods.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 5.7-7.2
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 19,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex crinita
FRINGED SEDGE

Native
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.

HABITAT: Moist to wet woods, thickets, marshes, ditches, streambanks.

CHARACTERISTICS:
Height: Up to 5.2 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4-7.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 720,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex intumescens
STAR SEDGE

Native
Bunch-type sedge; provides food and cover for wildlife.

HABITAT: Wet woods, meadows, swamps.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Shade
Drought Tolerance: No
pH: 4.8-6.9
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 40,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex frankii
FRANK'S SEDGE

Native
Very rhizomatous sedge; establishes quickly from seed; provides food for wildlife.

HABITAT: Swamps, wet woods, streambanks, ditches.

CHARACTERISTICS:
Height: Up to 2.6 ft.
Minimum Root Depth: 9 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 5.9-7.2
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 500,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex lupulina
HOP SEDGE

Native
Bunch-type sedge; ducks enjoy the large seed grains; provides food and cover for wildlife.

HABITAT: Swamps, wet woods.

CHARACTERISTICS:
Height: Up to 4.2 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 6.2-7
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 94,700
Seeding Rate: Up to 15% of a mix

Herbaceous Perennial



Carex lurida
LURID SEDGE

Native
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.

HABITAT: Swamps, bogs, wet meadows, wet woods.

CHARACTERISTICS:
Height: Up to 3.6 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.9-6.8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 250,000
Seeding Rate: Up to 20% of a mix

Herbaceous Perennial



Carex shortiana
SHORT'S SEDGE

Native
Bunch-type sedge; shortest sedge we grow; discovered by botanist Charles Wilkins Short; provides food and cover for wildlife.

HABITAT: Calcareous wet meadows, rich woods, swamps.

CHARACTERISTICS:
Height: Up to 2.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: No
pH: 4.7-6.9
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,000,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Carex pensylvanica
PENNSYLVANIA SEDGE

Native
Alternative stoloniferous sedge; tolerates sandy soils; ideal ground cover in a mature deciduous forest; may need prescribed fire to form a sod.

HABITAT: Open woods, wooded slopes.

CHARACTERISTICS:
Height: Up to 1.6 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Seeding Rate: Available as plants only

Herbaceous Perennial



Carex stipata
AWL SEDGE

Native
Bunch-type sedge; matures early in the season; provides food and cover for wildlife.

HABITAT: Wet meadows, swampy woods.

CHARACTERISTICS:
Height: Up to 4.2 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4.9-7.9
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 544,000
Seeding Rate: Up to 5% of a mix

Herbaceous Perennial



Carex scoparia
BLUNT BROOM SEDGE

Native
Bunch-type sedge; provides food and cover for songbirds, ruffed grouse chicks, ducks, and moose.

HABITAT: Swamps, wet meadows, moist open ground.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Shade
Drought Tolerance: No
pH: 4.6-6.9
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,344,000
Seeding Rate: Up to 15% of a mix

Herbaceous Perennial



Carex stricta
TUSsock SEDGE

Native
Tussock-forming sedge; provides habitat for turtles; source of food and cover for wildlife.

HABITAT: Swamps, streambanks, wet meadows.

CHARACTERISTICS:
Height: Up to 4.7 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 3.5-7
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,800,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Carex vulpinoidea
FOX SEDGE

Native
Bunch-type sedge; often the earliest sedge to establish from seed; provides food and cover for wildlife.

HABITAT: Moist meadows, marshes, ditches.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 6.8-8.9
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,297,000
Seeding Rate: Up to 35% of a mix

Herbaceous Perennial



Cinna arundinacea
WOOD REEDGRASS

Native
Bunch-type grass; excellent cover in forested wetlands; provides forage for wildlife.

HABITAT: Wet woods, swamps.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4-8.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 1,000,000
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Chasmanthium latifolium
RIVER OATS

Native
Decorative bunchgrass; adds variety and texture to wildflower mixes; great for riparian sites; provides food and cover for wildlife.

HABITAT: Riverbanks, alluvial woods.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
pH: 5-7
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 90,000
Seeding Rate: Up to 40% of a mix

Herbaceous Perennial



Cyperus esculentus
CHUFA

Native
Rhizomatous species; tubers are eaten by wildlife that dig them from the soil.

HABITAT: Moist ground of fields, meadows, lawns, gardens.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5-7
Bloom Period: Summer/Fall

Seeding Rate: Up to 5 lb per acre in a mix

Herbaceous Perennial



GOT A PLAN?

Talk to a pro today and get growing.

Whether you are tackling a small backyard project or a sprawling corporate campus, our pros are standing by ready to dig in with you. No question is too simple or too complex for our seasoned team of environmental enthusiasts.

Give us a ring Monday through Friday, 8 a.m. to 5 p.m. EST, or email us anytime.



TOLL-FREE
(800) 873-3321

PHONE
(814) 336-2404



E-MAIL
sales@ernstseed.com



Herbaceous Perennial

Dactylis glomerata
ORCHARDGRASS

Naturalized
Cool season bunchgrass with a dense root system; reliable grass for many grazing programs; provides excellent livestock forage.

HABITAT: Open fields, meadows, roadsides with well-drained medium-textured soils.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
pH: 5-7.5
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 427,000
Seeding Rate: Up to 10 lb per acre alone



Herbaceous Perennial

Elymus canadensis
CANADA WILDRYE

Native
Short-lived cool season bunchgrass; establishes quickly in disturbed areas; used for soil stabilization; provides food and cover for wildlife.

HABITAT: Alluvial thickets, streambanks, meadows; establishes best in well-drained soils.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
pH: 5-7.9
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 114,000
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

Deschampsia cespitosa
TUFTED HAIRGRASS

Native
Clump-forming grass.

HABITAT: Wet or boggy ground.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
pH: 3.5-7.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,100,000
Seeding Rate: Up to 1.3% of a mix; up to 0.25 lb per acre in a wildflower mix



Herbaceous Perennial

Elymus hystrix
BOTTLEBRUSH GRASS

Native
Short-lived cool season bunchgrass; used for soil stabilization; provides food and cover for wildlife.

HABITAT: Dry to mesic forests, woods.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Shade
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 75,000
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Herbaceous Annual

Echinochloa muricata
BARNYARDGRASS

Native
Excellent bunchgrass cover crop for moist and wet disturbed soils; Echinochloa crusgalli var. frumentacea (Japanese Millet) is a non-native substitute; provides food for songbirds and game birds.

HABITAT: Moist ground, alluvial shores.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Summer/Fall

Seeding Rate: Up to 10 lb per acre as a cover crop



Herbaceous Perennial

Elymus riparius
RIVERBANK WILDRYE

Native
Short-lived cool season bunchgrass; used for soil stabilization, often mixed with Virginia Wildrye; provides food and cover for wildlife.

HABITAT: Alluvial flats, meadows, streambanks, wet rich woods.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4.5-7.2
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 125,000
Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone



Elymus virginicus
VIRGINIA WILDRYE

Native

Short-lived cool season bunchgrass; used for soil stabilization and revegetation of wetlands, often found with Riverbank Wildrye; provides food and cover for wildlife.

HABITAT: Moist woods, meadows, riverbanks.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Shade

Drought Tolerance: Moderate

pH: 5-7

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 73,000

Seeding Rate: Up to 20% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Eragrostis spectabilis
PURPLE LOVEGRASS

Native

Low-growing, short-rhizomed bunchgrass; grows well in open areas; provides a short visual layer when used with little bluestem or sideoats grama; early fall color.

HABITAT: Sandy fields, pastures, roadsides, open woods, open areas; tolerates low fertility soils.

CHARACTERISTICS:

Height: Up to 2.6 ft.

Minimum Root Depth: 4 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 4-7.5

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 1,000,000

Seeding Rate: Up to 2% of a mix; up to 8 PLS lb per acre alone

Herbaceous Perennial



Eragrostis curvula
WEeping LOVEGRASS

Naturalized

Fast-establishing warm season bunchgrass; grows south of the Mason-Dixon Line; used for soil stabilization on steep slopes.

HABITAT: Sandy roadsides, fields.

CHARACTERISTICS:

Height: Up to 5.9 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 4.5-8.5

Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,482,000

Seeding Rate: Up to 5% of a mix; up to 3-5 PLS lb per acre alone

Herbaceous Perennial



Festuca arundinacea
TALL FESCUE

Naturalized

Cool season bunchgrass; used for mine reclamation, stabilizing grassed waterways, slopes, lawns, and recreation areas.

HABITAT: Low fertility acidic, clay, loamy, or sandy soils.

CHARACTERISTICS:

Height: Up to 6 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

pH: 5-9

Bloom Period: Spring/Summer

Approx Seeds Per Lb: 227,000

Seeding Rate: Up to 35% of a mix; up to 300 lb per acre alone

Herbaceous Perennial



Eragrostis hirsuta
BIGTOP LOVEGRASS

Native

Bunchgrass; attractive addition to upland meadows; early fall color.

HABITAT: Open disturbed habitats, clearings, roadsides, fields, open woods.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Shade Tolerance: Full Sun

Drought Tolerance: Unknown

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 1,000,000

Seeding Rate: Up to 2% of a mix; up to 8 PLS lb per acre alone

Herbaceous Perennial



Festuca ovina
SHEEP FESCUE

Naturalized

Fine-leaved cool season bunchgrass; adds texture to landscapes; used for reclamation, banks, and pastures.

HABITAT: Open woods, dry fields, roadsides.

CHARACTERISTICS:

Height: Up to 2.6 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Moderate

Drought Tolerance: High

pH: 5.5-7.5

Bloom Period: Spring/Summer

Approx Seeds Per Lb: 565,000

Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone

Herbaceous Perennial



Herbaceous Perennial

Festuca ovina var. duriuscula
HARD FESCUE

Naturalized
Fine-leaved cool season bunchgrass; used for reclamation and roadway revegetation where short vegetation is essential; good wildflower companion crop.

HABITAT: Low fertility well-drained soils.

CHARACTERISTICS:
Height: Up to 2.3 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 4.5-8.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 565,000
Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone



Herbaceous Perennial


Festuca rubra ssp. commutata
CHEWINGS FESCUE

Naturalized
Fine-leaved fescue; used for erosion control on slopes, waterways, and reclamation areas.

HABITAT: Well-drained acidic soils.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 500,000
Seeding Rate: Up to 35% of a mix; up to 220 lb per acre alone



Herbaceous Perennial

Festuca ovina var. glauca
BLUE FESCUE

Naturalized
Fine-leaved, ornamental cool season bunchgrass; used for ground cover and erosion control.

HABITAT: Open woods, roadsides, dry fields.

CHARACTERISTICS:
Height: Up to 2.3 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 550,000
Seeding Rate: Up to 35% of a mix; up to 25 lb per acre in a wildflower mix; up to 220 lb per acre alone



Herbaceous Perennial

Glyceria grandis
AMERICAN MANNAGRASS

Native
Decorative, wetland cool season bunchgrass; provides food for waterfowl, muskrats, and deer.

HABITAT: Swamps, marshes, wet meadows, shallow water, brooksides.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 659,000
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial

Festuca rubra
CREEPING RED FESCUE

Native
Sod-forming species; used for a no-mow cover and erosion control; provides habitat for wildlife.

HABITAT: Dry woods, roadsides, open ground.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
pH: 5-7.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 454,000
Seeding Rate: Up to 35% of a mix; up to 220 lb per acre alone



Herbaceous Perennial

Glyceria striata
FOWL MANNAGRASS

Native
Rhizomatous cool season bunchgrass; stays green through winter; provides food for waterfowl, muskrats, and deer.

HABITAT: Marshes, wet woods, swamps, bogs.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Minimum Root Depth: 4 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4-8
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,540,000
Seeding Rate: Up to 2% of a mix

Credit: Scott Rushe



Iris versicolor
BLUEFLAG

Native
Rhizomatous species; provides food for waterfowl, marsh birds, and muskrats.

HABITAT: Wet meadows, bogs, marshes.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Low
pH: 5-7
Bloom Period: Spring/Summer
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 18,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Koeleria macrantha
JUNEGRASS

Native
Cool season bunchgrass; provides good spring forage for livestock and deer; provides food for small mammals and upland game birds.

HABITAT: Open woods, dry soils.

CHARACTERISTICS:
Height: Up to 2 ft.
Minimum Root Depth: 20 in.
Shade Tolerance: Shade
Drought Tolerance: High
pH: 6-8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 2,315,000
Seeding Rate: Up to 1.5% of a mix

Herbaceous Perennial



Juncus effusus
SOFT RUSH

Native
Bunch-type grass with a wide geographic distribution; provides spawning grounds for bluegills in shallow water; source of food and cover for songbirds and waterfowl.

HABITAT: Swamps, moist fields, floodplains, shores, ditches.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 24 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.5-8.8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 45,359,000
Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Leersia oryzoides
RICE CUTGRASS

Native
Very rhizomatous warm season grass; creates a natural sediment trap; not recommended for use in residential settings as the vegetation can cause cuts to the skin (which gives the species its common name); provides food for ducks and habitat for invertebrates.

HABITAT: Marshes, bogs, wet meadows.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.1-8.8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 498,000
Seeding Rate: Up to 5% of a mix

Herbaceous Perennial



Juncus tenuis
PATH RUSH

Native
Bunch-type grass; tolerates foot traffic but not close mowing; used for nest material.

HABITAT: Moist to dry, often heavily compacted, soils of woods, fields, paths.

CHARACTERISTICS:
Height: Up to 2.6 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.5-7
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 29,000,000
Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Lolium multiflorum
ANNUAL RYEGRASS

Naturalized
Short-lived cool season bunchgrass; provides quick protection against soil, wind, and water erosion; used as a companion or cover crop where erosion is an immediate concern; frequently reseeds itself in disturbed areas.

HABITAT: Loose fertile to semi-fertile soils.

CHARACTERISTICS:
Height: Up to 2.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5-7.9
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 217,000
Seeding Rate: Up to 10% of a mix; up to 12 lb per acre as a companion crop with natives

Herbaceous Annual



Herbaceous Perennial

Onoclea sensibilis
SENSITIVE FERN

Native
Rhizomatous fern; provides shelter for salamanders and frogs.

HABITAT: Marshes, swamps, moist open woods, wet meadows.

CHARACTERISTICS:
Height: Up to 4.4 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 6,000,000
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial

Panicum clandestinum
DEERTONGUE

Native
Warm season bunchgrass with a fibrous root system; excellent for erosion control and revegetation of acid mine spoil and pipelines through wooded areas; good component of a streambank stabilization mix.

HABITAT: Moist soils of woodland edges and clearings.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 4-7.5
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 350,000
Seeding Rate: Up to 3% of a mix; up to 8 PLS lb per acre alone



Herbaceous Perennial

Panicum amarum
COASTAL PANICGRASS

Native
Warm season bunchgrass; used for the stabilization of coastal dunes, wind erosion control, and reclamation of gravel and mine areas; provides food and cover for wildlife.

HABITAT: Sandy shores, dune grasslands.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 5-7.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 325,000
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone



Herbaceous Annual

Panicum dichotomiflorum
SMOOTH PANICGRASS

Native
Warm season bunchgrass; used as a cover crop for wetland establishment; readily reseeds itself; plants and seeds provide food for ducks, deer, rabbits, and muskrats.

HABITAT: Moist soils, open woods, meadows.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
pH: 4.8-7
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 392,000
Seeding Rate: Up to 10% of a mix



Herbaceous Perennial

Panicum anceps
BEAKED PANICGRASS

Native
Bunchgrass; provides food and cover for wildlife; forage is of good value for cattle; seeds are eaten by upland birds.

HABITAT: Moist sandy soils of ditches, fields, savannas, low pinelands.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 225,000
Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

Panicum rigidulum
REDTOP PANICGRASS

Native
Bunchgrass; attractive seedheads and red foliage in late summer and early fall; common component of wetlands in the Southeast; palatable for livestock.

HABITAT: Wet soils of marshes, alluvial swamps, ditches, low woods.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5-7.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 797,000
Seeding Rate: Up to 35% of a mix



Panicum virgatum
SWITCHGRASS

Native

Rhizomatous warm season bunchgrass; used for biomass, soil stabilization on strip mine spoil and dikes, and in buffer strips for nutrient uptake; provides pasture and hay for cattle and sheep; source of food and cover for wildlife.

HABITAT: Brackish marshes, riverside prairies, open woods, prairies.

CHARACTERISTICS:

Height: Up to 6.1 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 4.5-8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 259,000
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone

Herbaceous Perennial



Poa trivialis
ROUGH BLUEGRASS

Naturalized

Cool season bunchgrass; good for use in detention basins; grows well in early spring; provides food and cover for wildlife.

HABITAT: Wet meadows, moist woods, roadsides.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4.8-7.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 2,500,000
Seeding Rate: Up to 25% of a mix; up to 160 lb per acre alone (lawns)

Herbaceous Perennial



Peltandra virginica
ARROW ARUM

Native

Bunch-type species; seeds must be stored cold and wet; seed pods containing numerous large seeds that ripen in the fall are a source of food for wood ducks, muskrats, and rails; foliage provides cover for aquatic mammals, wading birds, and waterfowl.

HABITAT: Swamps, stream or lake edges, tidal marshes.

CHARACTERISTICS:

Height: Up to 2.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: No
pH: 5-8.8
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 600
Seeding Rate: Up to 0.5 lb per 1,000 sq ft

Herbaceous Perennial



Pontederia cordata
PICKERELWEED

Native

Ornamental wetland bunchgrass; seeds must be stored cold and wet; provides food for wildlife.

HABITAT: Marshes, swampy edges of lakes and streams, along tidal shores.

CHARACTERISTICS:

Height: Up to 3.4 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
Pollinator Value: High
pH: 4.9-8.7
Bloom Period: Spring/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 5,000
Seeding Rate: Up to 10 lb per acre

Herbaceous Perennial



Poa palustris
FOWL BLUEGRASS

Native

Cool season bunchgrass; establishes quickly in wetlands and retention basins; provides food and cover for wildlife.

HABITAT: Wet meadows, damp soils.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.9-7.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 1,900,000
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre in a wet meadow mix; up to 160 lb per acre alone (lawns)

Herbaceous Perennial



Puccinellia distans
ALKALIGRASS

Naturalized

Bunchgrass; used for erosion control and along roadsides where salt runoff is prevalent.

HABITAT: Roadsides, disturbed sites.

CHARACTERISTICS:

Height: Up to 2.3 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Bloom Period: Spring/Fall

Approx Seeds Per Lb: 1,200,000
Seeding Rate: Up to 20% of a mix

Herbaceous Perennial



Sagittaria latifolia
DUCK POTATO

Native
Ornamental wetland species; seeds and large tubers provide food for waterfowl, songbirds, wading birds, muskrats, and beavers.

HABITAT: Swamps, wet shores, shallow water of ponds and streams.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
Pollinator Value: Low
pH: 4.7-8.9
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 67,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Scirpus cyperinus
WOOLGRASS

Native
Bunch-type species; stands tall in marshes; provides food and cover for waterfowl and muskrats.

HABITAT: Moist meadows, marshes, swamps, shores, ditches.

CHARACTERISTICS:
Height: Up to 5.6 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.8-7.2
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 36,000,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Schizachyrium scoparium
LITTLE BLUESTEM

Native
Long-lived bunchgrass; dense roots can grow to 8' deep; good for upland meadows where sight lines are important; used for erosion control on droughty sites; provides summer forage for livestock; source of food and cover for wildlife; grows with open exposed surfaces that host ground nesting pollinators.

HABITAT: Old fields, roadsides, riverside prairies, open woods, slopes, meadows.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 5-8.4
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 241,000
Seeding Rate: Up to 30% of a mix; up to 10 PLS lb per acre alone

Herbaceous Perennial



Scirpus expansus
WOOD BULRUSH

Native
Very rhizomatous species; provides cover for wildlife.

HABITAT: Marshes, wet meadows, swamps, swales.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 10,000,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Scirpus atrovirens
GREEN BULRUSH

Native
Bunch-type species; provides cover for wildlife in wet areas.

HABITAT: Moist meadows, marshes, floodplain forests, ditches; tolerates clay and silt soils found in wet areas.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4-8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 11,300,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Scirpus hattorianus
NORTHERN BULRUSH

Native
Clump-forming species.

HABITAT: Swamps, bogs, moist meadows, riverbanks, ditches.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 36,000,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Scirpus pendulus
BROWN BULRUSH

Native
Bunch-type species; used for wildlife habitat.

HABITAT: Wet meadows, marshes, swales, ditches.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
pH: 4.9-7
Bloom Period: Spring

Approx Seeds Per Lb: 30,000,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Sorghastrum nutans
INDIANGRASS

Native
Attractive warm season bunchgrass; good for erosion control, landscaping, and roadside beautification; provides food and cover for wildlife.

HABITAT: Riverside prairies, moist or dry fields, open woods, roadsides, serpentine barrens; grows best in deep well-drained soils.

CHARACTERISTICS:

Height: Up to 8.1 ft.
Minimum Root Depth: 24 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 4.8-8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 175,000
Seeding Rate: Up to 30% of a mix; up to 8 PLS lb per acre alone

Herbaceous Perennial



Scirpus validus
SOFTSTEM BULRUSH

Native
Rhizomatous species; provides spawning grounds for fish in shallow water; source of food for waterfowl and muskrats.

HABITAT: Swamps, wet ditches, mud flats, pond and lake margins.

CHARACTERISTICS:

Height: Up to 9.8 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
pH: 5.4-7.5
Bloom Period: Summer/Fall
Flower Color(s): Red

Approx Seeds Per Lb: 496,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Sparganium americanum
EASTERN BUR REED

Native
Rhizomatous, emergent aquatic plant; provides food for waterfowl, muskrats, and beavers.

HABITAT: Muddy shores, shallow water rivers, streams, swamps, ponds.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: No
pH: 4.9-7.3
Bloom Period: Spring/Fall
Flower Color(s): Green

Approx Seeds Per Lb: 50,000
Seeding Rate: Up to 20% of a mix

Herbaceous Perennial



Sisyrrinchium angustifolium
NARROWLEAF BLUE EYED GRASS

Native
Early season food source for pollinators.

HABITAT: Damp woods, floodplains, grassy places, meadows.

CHARACTERISTICS:

Height: Up to 1.6 ft.
Minimum Root Depth: 4 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Medium
pH: 5-7
Bloom Period: Spring/Summer
Flower Color(s): Blue

Approx Seeds Per Lb: 757,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Sparganium eurycarpum
GIANT BUR REED

Native
Rhizomatous, emergent aquatic plant; provides food and cover for waterfowl, pheasants, muskrats, and beavers.

HABITAT: Bogs, swamps, lake margins, ditches, swampy meadows.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: No
pH: 5-8.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 1,500
Seeding Rate: Up to 20% of a mix

Herbaceous Perennial



Herbaceous Perennial

Spartina pectinata
PRAIRIE CORDGRASS

Native
Aggressive, rhizomatous warm season grass; root system provides erosion control on streambanks; source of cover for waterfowl, songbirds, and small mammals.

HABITAT: Marshes, wet prairies, shores, riverside prairies.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 6-8.5
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 639,000
Seeding Rate: Up to 5% of a mix; up to 8 PLS lb per acre alone



Herbaceous Perennial

Sporobolus heterolepis
PRAIRIE DROPSEED

Native
Decorative, fine-textured warm season bunchgrass; may be difficult to germinate; provides food and cover for wildlife.

HABITAT: Dry open ground.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
pH: 6-7.2
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 1,200,000
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

Sporobolus asper
ROUGH DROPSEED

Native
Warm season bunchgrass; survives in very poor soil conditions with little organic matter.

HABITAT: Dry sandy soils.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 5.5-7
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 760,000
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial

Tradescantia ohiensis
OHIO SPIDERWORT

Native
Attractive clump-forming species; blooms open during the morning hours from spring through summer; flowers close in warm temperatures.

HABITAT: Meadows, prairies, thickets, dry rocky woodlands, floodplain forests.

CHARACTERISTICS:
Height: Up to 4.1 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Spring/Summer
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 128,000
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Sporobolus cryptandrus
SAND DROPSEED

Native
Early emerging warm season bunchgrass; provides food and cover for wildlife.

HABITAT: Dry sandy soils.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 6.6-8
Bloom Period: Summer/Fall

Approx Seeds Per Lb: 5,600,000
Seeding Rate: Up to 0.1% of a mix



Herbaceous Perennial

Tradescantia subaspera
ZIGZAG SPIDERWORT

Native
Attractive clump-forming species; flowers close in warm temperatures.

HABITAT: Mesic to dry forests and woodlands.

CHARACTERISTICS:
Height: Up to 4.1 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 128,000
Seeding Rate: Up to 2% of a mix



Tradescantia virginiana
VIRGINIA SPIDERWORT

Native

Attractive, late spring clump-forming species; more compact than Ohio Spiderwort; flowers close in warm temperatures.

HABITAT: Prairies, mesic to dry upland forests, floodplain forests.

CHARACTERISTICS:

Height: Up to 2.9 ft.

Minimum Root Depth: 4 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

Pollinator Value: High

pH: 4-8

Bloom Period: Spring/Summer

Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 175,000

Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Tripsacum dactyloides
EASTERN GAMAGRASS

Native

Warm season bunchgrass; plant in the fall as a dormant seeding with germination occurring in the spring; requires great patience as the seeds have high levels of dormancy; outstanding forage producer; provides food and cover for wildlife.

HABITAT: Riverside prairies, meadows, swamps, wet shores, open fields.

CHARACTERISTICS:

Height: Up to 9.8 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 5.1-7.5

Bloom Period: Spring/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 7,000

Seeding Rate: Up to 50% of a mix; up to 20 PLS lb per acre alone

Herbaceous Perennial



Tridens flavus
PURPLETOP

Native

Warm season bunchgrass; adds late season color to native landscapes; used for soil stabilization; provides food and cover for wildlife.

HABITAT: Meadows, old fields, open woods, roadsides; tolerates low quality roadside and field soils.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 4.5-6.5

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 465,000

Seeding Rate: Up to 15% of a mix; up to 10 PLS lb per acre alone drill seeded; up to 20 PLS lb per acre alone broadcast seeded

Herbaceous Perennial



Zizania aquatica
WILDRICE

Native

Bunchgrass; seeds must be kept moist and require clean still water to grow; provides food for wood ducks, black ducks, and muskrats.

HABITAT: Tidal and non-tidal marshes that are 1" deep or more.

CHARACTERISTICS:

Height: Up to 9.8 ft.

Minimum Root Depth: 6 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

pH: 6.4-7.4

Bloom Period: Summer/Fall

Approx Seeds Per Lb: 11,000

Seeding Rate: Up to 10 lb per acre when used with a mix; up to 40 lb per acre alone

Herbaceous Annual

ORDER TODAY

Placing your order should be the easy part.

You're busy, so ordering should be the easiest part of the project. We accept orders by phone, e-mail, or by mail.

ORDER MINIMUMS

- ▶ \$25.00 order minimum before freight and tax.
- ▶ Mixes must be ordered in whole pound increments.
- ▶ 5% custom mix charge.

PAYMENT

Visa, MasterCard, Discover, and American Express are accepted forms of payment. Checks and money orders are also acceptable with placement of order. Sorry, Ernst cannot process COD orders.



TOLL-FREE
(800) 873-3321

PHONE
(814) 336-2404



E-MAIL
sales@ernstseed.com



Herbaceous Perennial

Achillea millefolium
COMMON YARROW

Naturalized
Species with a long bloom period.
HABITAT: Roadsides, fields, waste areas.
CHARACTERISTICS:
Height: Up to 3.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
pH: 6-8
Bloom Period: Spring/Fall
Flower Color(s): White
Approx Seeds Per Lb: 2,852,000
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

Amsonia ciliata
FRINGED BLUESTAR

Native
Clump-forming species; provides food for native pollinators.
HABITAT: Sandhills, sandy woodlands.
CHARACTERISTICS:
Height: Up to 2.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Bloom Period: Spring
Flower Color(s): Blue
Approx Seeds Per Lb: 45,400
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

Agastache foeniculum
ANISE (LAVENDER) HYSSOP

Native
Provides food for pollinators.
HABITAT: Dry upland woods and prairies.
CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Shade
Drought Tolerance: Low
Bloom Period: Summer/Fall
Flower Color(s): Blue
Approx Seeds Per Lb: 1,540,000
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

Anemone canadensis
CANADIAN ANEMONE

Native
Very attractive rhizomatous species.
HABITAT: Sandy shores, thickets, damp prairies, wet meadows.
CHARACTERISTICS:
Height: Up to 2.6 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): White
Approx Seeds Per Lb: 131,660
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Agrimonia parviflora
SMALL FLOWERED AGRIMONY

Native
Adds structure and texture to wetlands.
HABITAT: Bogs, moist woods, thickets; tolerates poor soils.
CHARACTERISTICS:
Height: Up to 5.9 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 6-8
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 288,000
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Anemone virginiana
THIMBLEWEED

Native
Excellent for moderately shaded areas.
HABITAT: Dry open woods, slopes, thickets.
CHARACTERISTICS:
Height: Up to 3.8 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Low
Bloom Period: Spring/Summer
Flower Color(s): Green
Approx Seeds Per Lb: 448,000
Seeding Rate: Up to 1% of a mix



Apocynum cannabinum
INDIAN HEMP

Native

Spreads by underground roots; used by Native Americans to make rope; may be allelopathic; stem fiber is used by songbirds and orioles to build nests; flowers are attractive to butterflies.

HABITAT: Woods, old fields, sandy flats, limestone bluffs, open ground.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

Pollinator Value: Very High

pH: 4.5-7

Bloom Period: Spring/Fall

Approx Seeds Per Lb: 500,000

Seeding Rate: Up to 0.3% of a mix

Herbaceous Perennial



Asclepias syriaca
COMMON MILKWEED

Native

Decorative species with a fragrance resembling a lilac; spreads from underground roots; essential food source for monarch butterfly caterpillars and other beneficial insects.

HABITAT: Fields, roadsides, open ground.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Summer

Flower Color(s): Purple

Approx Seeds Per Lb: 70,000

Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Aquilegia canadensis
EASTERN COLUMBINE

Native

One of the first flowers to bloom in the spring; nectar source for hummingbirds.

HABITAT: Cliffs, rocky slopes, dry woods; usually calcareous.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Moderate

Drought Tolerance: Unknown

Pollinator Value: Low

pH: 5-8

Bloom Period: Spring/Summer

Flower Color(s): Red

Approx Seeds Per Lb: 504,000

Seeding Rate: Up to 1.3% of a mix

Herbaceous Perennial



Asclepias tuberosa
BUTTERFLY MILKWEED

Native

Showy clump-forming species with tuberous roots; essential food source for monarch butterfly caterpillars.

HABITAT: Dry woods, abandoned fields, roadsides, shale barrens; grows best in well-drained soils.

CHARACTERISTICS:

Height: Up to 2.9 ft.

Minimum Root Depth: 16 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

Pollinator Value: Very High

pH: 4.8-6.8

Bloom Period: Spring/Fall

Flower Color(s): Orange

Approx Seeds Per Lb: 70,000

Seeding Rate: Up to 2.5% of a mix

Herbaceous Perennial



Asclepias incarnata
SWAMP MILKWEED

Native

Decorative rhizomatous species with a fragrance of bubble gum; essential food source for monarch butterfly caterpillars; we observe more chrysalis on this milkweed species than on any other in our production fields.

HABITAT: Swamps, floodplains, wet meadows.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 18 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

Pollinator Value: Very High

pH: 5-8

Bloom Period: Summer/Fall

Flower Color(s): Pink

Approx Seeds Per Lb: 153,760

Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Aster divaricatus
WHITE WOOD ASTER

Native

Rhizomatous species; widely distributed in wooded areas; provides food and habitat for wildlife.

HABITAT: Dry woods.

CHARACTERISTICS:

Height: Up to 3.3 ft.

Shade Tolerance: Moderate

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Summer/Fall

Flower Color(s): White

Approx Seeds Per Lb: 670,000

Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster laevis
SMOOTH BLUE ASTER

Native
Provides attractive late summer color in meadows; sought by deer for browse.

HABITAT: Dry woods, rocky ledges, roadsides.

CHARACTERISTICS:
Height: Up to 6.3 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5.8-7.8
Bloom Period: Summer/Fall
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 1,014,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster lowrieanus
LOWRIE'S BLUE WOOD ASTER

Native
Source of food for pollinators.

HABITAT: Dry to mesic woodlands.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Moderate
Drought Tolerance: No
pH: 5.7-7.5
Bloom Period: Summer/Fall
Flower Color(s): Blue

Approx Seeds Per Lb: 2,000,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Aster lanceolatus
LANCE LEAVED ASTER

Native
Rhizomatous species; branches provide texture to landscapes.

HABITAT: Open woods, old fields, roadsides, moist low places.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5.8-7.4
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 700,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster macrophyllus
BIGLEAF ASTER

Native
Showy rhizomatous species; source of color along wooded borders; provides habitat for wildlife.

HABITAT: Woods, rocky slopes, woodland edges; grows best in fertile soils.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Shade
Drought Tolerance: Low
Pollinator Value: Very High
pH: 4.9-6.9
Bloom Period: Summer/Fall
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 800,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster lateriflorus
CALICO ASTER

Native
Attractive clump-forming species; provides food and cover for wildlife.

HABITAT: Dry open places, open woods.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5.2-7.5
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 800,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster novae-angliae
NEW ENGLAND ASTER

Native
Brilliant flowers stand out from a distance; provides food and cover for wildlife.

HABITAT: Fields, roadsides, moist meadows.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 1,100,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster oblongifolius
AROMATIC ASTER

Native
Longest blooming of our native asters, lasting up to and sometimes beyond two months; small oblong leaves have a lemony scent.

HABITAT: Dry open places, calcareous hillsides, cliffs, bluffs.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 816,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster puniceus
PURPLESTEM ASTER

Native
Attractive rhizomatous species; adds color to FACW and OBL meadows; provides food for deer.

HABITAT: Swamps, wet meadows, riverbanks, moist roadsides.

CHARACTERISTICS:
Height: Up to 8.1 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
Pollinator Value: Very High
pH: 4.5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 700,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster pilosus
HEATH ASTER

Native
Aggressive rhizomatous species; establishes quickly on disturbed sites and rights-of-way.

HABITAT: Dry fields, open woods, vacant lots, roadsides.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5.4-7
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 700,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster sagittifolius
ARROWLEAF ASTER

Native
Rhizomatous species.

HABITAT: Woodland edges, streambanks, open areas, roadside slopes.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 1,487,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Aster prenanthoides
ZIGZAG ASTER

Native
Rhizomatous species; tolerates urban habitats.

HABITAT: Streambanks, wet meadows, low woods.

CHARACTERISTICS:
Height: Up to 5.5 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5.5-7.2
Bloom Period: Summer/Fall
Flower Color(s): Blue, Purple

Approx Seeds Per Lb: 700,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster spectabilis
SHOWY ASTER

Native
Attractive rhizomatous species; one of the prettiest asters to add to landscapes.

HABITAT: Woodland borders, pine barrens, clearings, roadsides, dry sandy soils.

CHARACTERISTICS:
Height: Up to 2.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Purple, Blue

Approx Seeds Per Lb: 630,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Aster umbellatus
**FLAT TOPPED WHITE
ASTER**

Native
Decorative rhizomatous component of wet meadows.

HABITAT: Floodplains, swamps, moist woods, moist fields.

CHARACTERISTICS:
Height: Up to 8.1 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 1,072,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Baptisia pendula
LARGELEAF WILD INDIGO

Native
Rhizomatous early spring legume.

HABITAT: Flatwoods, open woods, clearings.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 25,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial Legume



Baptisia alba
WHITE WILD INDIGO

Native
Slow-developing legume; lasts for many years in low fertility soils.

HABITAT: Open upland woods, prairies.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 25,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial Legume



Baptisia tinctoria
YELLOW FALSE INDIGO

Native
Essential food source for frosted elfin butterfly caterpillars.

HABITAT: Dry sandy soils, clearings, dry open woods.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
Pollinator Value: Medium
pH: 5.8-7
Bloom Period: Spring/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 300,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial Legume



Baptisia australis
BLUE FALSE INDIGO

Native
Historically used as a dye; may be used as a specimen plant.

HABITAT: Open woods, riverbanks, sandy floodplains.

CHARACTERISTICS:
Height: Up to 5.2 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): Purple

Approx Seeds Per Lb: 22,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial Legume



Bidens aristosa
**SHOWY TICKSEED
SUNFLOWER (BUR
MARIGOLD)**

Native
Reseeding annual in disturbed soils; provides aggressive first-year cover in wet meadows; seeds are eaten by wildlife.

HABITAT: Meadows, fields, roadsides, ditches.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Medium
pH: 5-7
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 130,000
Seeding Rate: Up to 2% of a mix

Herbaceous Annual



Bidens cernua
NODDING BUR MARIGOLD

Native

Reseeding annual persistent in disturbed saturated soils; not for use in residential settings as seeds attach themselves to clothing; provides food and cover for wildlife.

HABITAT: Swamps, wet shores, ditches.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Medium
pH: 5.1-7
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 43,324
Seeding Rate: Up to 2% of a mix

Herbaceous Annual



Centaurea cyanus
**CORNFLOWER
(BACHELOR'S BUTTON)**

Naturalized

Blooms in early spring if sown in the fall.

HABITAT: Meadows, flower beds.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Fall
Flower Color(s): Blue

Approx Seeds Per Lb: 90,000
Seeding Rate: Up to 11% of a mix

Herbaceous Annual



Bidens frondosa
BEGGARTICK

Native

Reseeding annual in disturbed wetlands; not for use in residential settings as seeds attach themselves to clothing; provides food and cover for wildlife and ducks.

HABITAT: Moist open ground, streambanks, roadsides.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Medium
pH: 5.2-7.2
Bloom Period: Summer/Fall
Flower Color(s): Yellow, Orange

Approx Seeds Per Lb: 195,300
Seeding Rate: Up to 1.3% of a mix

Herbaceous Annual



Chamaecrista fasciculata
PARTRIDGE PEA

Native

Bunch-type, readily reseeding annual in disturbed upland sites; foliage is nutritious but can be poisonous and should be considered potentially dangerous to cattle; fruit and seeds can irritate the digestive tract of livestock; seeds are a source of food for quail.

HABITAT: Riverbanks, sandy soils, clearings, roadsides.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
Pollinator Value: High
pH: 5.5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 65,000
Seeding Rate: Up to 3% of a mix

Herbaceous Annual Legume



Blephilia ciliata
DOWNY PAGODA PLANT

Native

Our earliest blooming mint; a beautiful addition to landscapes.

HABITAT: Dry woods, fields.

CHARACTERISTICS:

Height: Up to 3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: High
Bloom Period: Spring/Summer
Flower Color(s): Purple, White

Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Chamaecrista nictitans
SENSITIVE PEA

Native

Tap-rooted legume; can help to increase nitrogen availability in some soils; fruit and seeds can irritate the digestive tract of livestock; seeds provide food for quail, doves, and turkey; attracts parasitoid wasps that control plant-eating insects.

HABITAT: Fields, roadsides, clearings, near rivers, woodlands.

CHARACTERISTICS:

Height: Up to 1.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 206,570
Seeding Rate: Up to 1% of a mix

Herbaceous Annual Legume



Cheiranthus allionii
WALLFLOWER

Naturalized
Decorative early blooming component in wildflower mixes.

HABITAT: Fields, meadows.
CHARACTERISTICS:
Height: Up to 1.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring
Flower Color(s): Yellow, Orange, Red

Approx Seeds Per Lb: 345,000
Seeding Rate: Up to 3% of a mix

Herbaceous Biennial/Perennial



Cichorium intybus
BLUE CHICORY

Naturalized
Attractive bunch-type species; persistent on roadsides and in compacted soils; blooms last all day.

HABITAT: Fields, roadsides, waste ground.
CHARACTERISTICS:
Height: Up to 5.5 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 6-7.5
Bloom Period: Spring/Fall
Flower Color(s): Blue

Approx Seeds Per Lb: 426,400
Seeding Rate: Up to 2.5% of a mix

Herbaceous Perennial



Chrysanthemum maximum
SHASTA DAISY

Naturalized
Attractive component of naturalized meadow mixes.

HABITAT: Gardens, meadows.
CHARACTERISTICS:
Height: Up to 3 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 400,000
Seeding Rate: Up to 2.5% of a mix

Herbaceous Perennial



Clematis virginiana
VIRGIN'S BOWER

Native
Hardy vine to include in wetland margins; attractive when in bloom and when seed is ripening.

HABITAT: Thickets, streambanks, low woods.
CHARACTERISTICS:
Height: Up to 16.3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: Medium
pH: 5-6.8
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 192,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial Vine



FIND IT FAST

Try our Seed Finder Tool to narrow the search.

Finding the right seeds you need for your project can be frustrating. Save time by using the advanced search and filtering options of our Seed Finder Tool.

Try it now by scanning this QR code and making your selections. If you have any problems, feel free to reach out to our sales department for guidance.





Coreopsis basalis
GOLDENMANE TICKSEED

Native
Adds early and mid-season color to southeastern landscapes.

HABITAT: Fields and roadsides in sandy soils.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 500,000
Seeding Rate: Up to 2% of a mix

Herbaceous Annual



Coreopsis tinctoria
PLAINS COREOPSIS

Native
Showy annual; produces flowers in a short period of time on low fertility sites.

HABITAT: Fields, meadows, roadsides, occasionally escaping to yards.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Medium
pH: 5.2-7.8
Bloom Period: Spring/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 3,222,000
Seeding Rate: Up to 1% of a mix

Herbaceous Annual



Coreopsis grandiflora
LARGEFLOWER TICKSEED

Native
Showy component in wildflower meadows.

HABITAT: Roadsides, upland woods.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Shade Tolerance: Full Sun
Drought Tolerance: High
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 200,000
Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Coreopsis tripteris
TALL COREOPSIS

Native
Long-lived species; very tolerant of competition.

HABITAT: Old fields, thickets, woodland edges, roadsides, moist low places.

CHARACTERISTICS:

Height: Up to 9.8 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 200,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Coreopsis lanceolata
LANCELEAF COREOPSIS

Native
Popular for wildflower meadows and along roadsides.

HABITAT: Soils, thickets, fields, clearings, roadsides.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
Pollinator Value: Medium
pH: 6-7
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 221,000
Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Coronilla varia
CROWNVETCH

Naturalized
Rhizomatous legume; good for controlling erosion on steep, dry rocky slopes; seeds are persistent in soils; can be aggressive in natural areas.

HABITAT: Rocky slopes, shallow, well-drained low fertility soils.

CHARACTERISTICS:

Height: Up to 3.9 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: High
pH: 5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Pink

Approx Seeds Per Lb: 140,000
Seeding Rate: Up to 50% of a mix; 8-15 lb per acre

Herbaceous Perennial Legume



Herbaceous Annual

Cosmos bipinnatus
COSMOS

Naturalized
Robust with intense color; ideal for showy roadside plantings; attracts butterflies.

HABITAT: Disturbed sites, fields, roadsides.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Bloom Period: Summer/Fall
Flower Color(s): Mixed

Approx Seeds Per Lb: 72,000
Seeding Rate: Up to 13% of a mix



Herbaceous Perennial Legume

Dalea purpurea
PURPLE PRAIRIE CLOVER

Native
Fixes nitrogen in prairie and meadow soils; excellent forage for wildlife.

HABITAT: Dry prairies, open glades.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
pH: 6-8
Bloom Period: Spring/Summer
Flower Color(s): Purple

Approx Seeds Per Lb: 300,000
Seeding Rate: Up to 1% of a mix



Herbaceous Annual

Cosmos sulphureus
SULPHUR COSMOS

Naturalized
Attractive summer color lasts into the fall.

HABITAT: Gardens, roadsides, other disturbed habitats.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Full Sun
Drought Tolerance: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 60,000
Seeding Rate: Up to 15% of a mix



Herbaceous Biennial

Daucus carota
QUEEN ANNE'S LACE

Naturalized
Persistent in meadows with poor soils; attracts butterflies.

HABITAT: Roadsides, old fields, gardens, open ground.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Fall
Flower Color(s): White

Approx Seeds Per Lb: 454,000
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial Legume

Dalea candida
WHITE PRAIRIE CLOVER

Native
Supplies nitrogen to meadow soils; produces nutritious forage for deer; seeds are eaten by birds.

HABITAT: Dry prairies, dry upland woods.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: No
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 278,000
Seeding Rate: Up to 1% of a mix



Herbaceous Annual

Delphinium ajacis
ROCKET LARKSPUR

Naturalized
Attractive tap-rooted species.

HABITAT: Gardens, roadsides, other disturbed habitats.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: No
Bloom Period: Spring/Fall
Flower Color(s): Blue, White, Pink

Approx Seeds Per Lb: 140,000
Seeding Rate: Up to 5% of a mix



Desmanthus illinoensis
**ILLINOIS
BUNDLEFLOWER**

Native

Legume with a deep taproot; fixes nitrogen in prairie and meadow soils that can be used by plants.

HABITAT: Prairies, roadsides, meadows, riverbanks.

CHARACTERISTICS:

Height: Up to 7.8 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

pH: 5-8

Bloom Period: Summer

Flower Color(s): White

Approx Seeds Per Lb: 85,000

Seeding Rate: Up to 1% of a mix

Herbaceous Perennial Legume



Dianthus barbatus
SWEETWILLIAM

Naturalized

Food source for hummingbirds and butterflies.

HABITAT: Gardens, roadsides, other disturbed habitats; grows best in fertile, moist well-drained soils.

CHARACTERISTICS:

Height: Up to 2 ft.

Shade Tolerance: Moderate

Drought Tolerance: Unknown

Bloom Period: Spring/Summer

Flower Color(s): Red, Pink, White

Approx Seeds Per Lb: 440,000

Seeding Rate: Up to 2.5% of a mix

Herbaceous Perennial



Desmodium canadense
SHOWY TICKTREFOIL

Native

Nitrogen-fixing legume; not for use in residential settings as seeds attach themselves to clothing; seeds provide food for ground birds and small animals; attracts native pollinators.

HABITAT: Open woods, meadows, thickets, riverbanks.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Shade Tolerance: Moderate

Drought Tolerance: Unknown

Bloom Period: Summer/Fall

Flower Color(s): Purple

Approx Seeds Per Lb: 72,500

Seeding Rate: Up to 2% of a mix

Herbaceous Perennial Legume



Echinacea pallida
**PALE PURPLE
CONEFLOWER**

Native

Attractive in wildflower meadows.

HABITAT: Dry open spaces.

CHARACTERISTICS:

Height: Up to 3.3 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

pH: 6.5-7.2

Bloom Period: Summer

Flower Color(s): Pink, Purple

Approx Seeds Per Lb: 106,000

Seeding Rate: Up to 7% of a mix

Herbaceous Perennial



Desmodium paniculatum
**PANICLELEAF
TICKTREFOIL**

Native

Nitrogen-fixing legume; not for use in residential settings as seeds attach themselves to clothing; seeds provide food for wildlife; attracts native pollinators.

HABITAT: Dry woods, fields.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Minimum Root Depth: 6 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

Pollinator Value: Low

pH: 6-7

Bloom Period: Summer/Fall

Flower Color(s): Purple

Approx Seeds Per Lb: 200,000

Seeding Rate: Up to 1% of a mix

Herbaceous Perennial Legume



Echinacea purpurea
PURPLE CONEFLOWER

Native

Source of herbal remedy; attractive species in meadows and along roadsides.

HABITAT: Open woods, open meadows, prairies, roadsides; grows best in moist well-drained soils.

CHARACTERISTICS:

Height: Up to 5.9 ft.

Minimum Root Depth: 24 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

Pollinator Value: High

pH: 6.5-7.2

Bloom Period: Summer/Fall

Flower Color(s): Purple

Approx Seeds Per Lb: 115,664

Seeding Rate: Up to 7% of a mix

Herbaceous Perennial



Herbaceous Perennial

Eryngium yuccifolium
RATTLESNAKE MASTER

Native
Creates visual texture in native meadows; essential food source for the rare rattlesnake master borer moth.

HABITAT: Moist woods, moist or dry sandy soils, meadows, barrens.

CHARACTERISTICS:
Height: Up to 5.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer
Flower Color(s): White

Approx Seeds Per Lb: 178,000
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Eupatorium maculatum
SPOTTED JOE PYE WEED

Native
Source of food for pollinators.

HABITAT: Floodplains, swamps, alluvial thickets.

CHARACTERISTICS:
Height: Up to 8.1 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 1,440,000
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

Eupatorium coelestinum
MISTFLOWER

Native
Rhizomatous, low-growing, late fall flower in wetland margins.

HABITAT: Floodplain forests, old fields, meadows, streambanks.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5.5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Blue

Approx Seeds Per Lb: 1,500,000
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

Eupatorium perfoliatum
BONESET

Native
Beautiful, hardy component of wetlands; seeds are eaten by swamp sparrows.

HABITAT: Floodplains, swamps, bogs, streambanks, wet meadows.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: No
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 2,880,000
Seeding Rate: Up to 0.5% of a mix



Herbaceous Perennial

Eupatorium fistulosum
JOE PYE WEED

Native
Showy, rhizomatous, hollow-stemmed Joe Pye Weed.

HABITAT: Floodplains, meadows, moist thickets, roadsides.

CHARACTERISTICS:
Height: Up to 9.8 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: High
pH: 4.5-7
Bloom Period: Summer/Fall
Flower Color(s): Pink, Purple

Approx Seeds Per Lb: 2,000,000
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

Eupatorium purpureum
PURPLE NODE JOE PYE WEED

Native
Clump-forming species; stems have a sweet scent resembling vanilla when bruised.

HABITAT: Thickets, open woods.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Pink, Purple

Approx Seeds Per Lb: 672,000
Seeding Rate: Up to 1% of a mix



Euthamia graminifolia
GRASSLEAF GOLDENROD

Native
Rhizomatous species; provides food and cover for wildlife.

HABITAT: Wet meadows, riparian areas; tolerates poor soils.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 5,600,000
Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Gaura biennis
BIENNIAL BEEBLOSSOM

Native
Adds diversity to meadow mixes.

HABITAT: Moist meadows, streambanks, floodplains, roadside thickets.

CHARACTERISTICS:

Height: Up to 6.5 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Bloom Period: Summer/Fall
Flower Color(s): Pink, White

Approx Seeds Per Lb: 43,130
Seeding Rate: Up to 2% of a mix

Herbaceous Biennial



Gaillardia aristata
PERENNIAL GAILLARDIA (BLANKETFLOWER)

Native
Attractive bunch-type, daisy-like flower for meadows and along roadsides.

HABITAT: Plains, prairies, meadows, along roadsides.

CHARACTERISTICS:

Height: Up to 2.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.5-7.9
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 186,436
Seeding Rate: Up to 10% of a mix

Herbaceous Perennial



Geum canadense
WHITE AVENS

Native
Small seedheads add texture to landscapes after blooming.

HABITAT: Dry or moist woods, roadsides.

CHARACTERISTICS:

Height: Up to 3.6 ft.
Minimum Root Depth: 4 in.
Shade Tolerance: Shade
Drought Tolerance: Low
Pollinator Value: Medium
pH: 4.5-7.5
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 400,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Gaillardia pulchella
ANNUAL GAILLARDIA (INDIAN BLANKET)

Native
Showy bunch-type species in annual wildflower beds.

HABITAT: Dry sandy places, meadows, open areas.

CHARACTERISTICS:

Height: Up to 2 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 7-8.5
Bloom Period: Spring/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 238,144
Seeding Rate: Up to 10% of a mix

Herbaceous Annual/Biennial/Perennial



Gypsophila elegans
ANNUAL BABY'S BREATH

Naturalized
Fine-textured, fast-growing species; blooms 60 days after seeding.

HABITAT: Gardens, meadows.

CHARACTERISTICS:

Height: Up to 2 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Bloom Period: Summer
Flower Color(s): White

Approx Seeds Per Lb: 375,000
Seeding Rate: Up to 2.5% of a mix


Herbaceous Annual



Helenium autumnale
COMMON SNEEZEWEED

Native
Attractive late season bloomer.
HABITAT: Swamps, moist riverbanks, alluvial thickets, wet fields.
CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
Pollinator Value: High
pH: 4-7.5
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 1,464,500
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Helianthus annuus
COMMON SUNFLOWER

Native
Reseeding annual; provides food for songbirds.
HABITAT: Meadows, roadsides.
CHARACTERISTICS:
Height: Up to 11.4 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.5-7.8
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 46,900
Seeding Rate: Up to 10% of a mix

Herbaceous Annual



Helenium flexuosum
PURPLEHEAD SNEEZEWEED

Native
Provides summer color to moist soils.
HABITAT: Wet meadows, riverbanks, moist fields, roadsides.
CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: High
pH: 4.5-7.5
Bloom Period: Spring/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 2,000,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Helianthus maximilianii
MAXIMILIAN'S SUNFLOWER

Native
Clump-forming species; tall shielding growth provides food and cover for birds.
HABITAT: Prairies, old fields, railroad tracks, urban open ground.
CHARACTERISTICS:
Height: Up to 9.8 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 6-8
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 196,300
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Helianthus angustifolius
NARROWLEAF SUNFLOWER

Native
Clump-forming species; latest blooming sunflower we carry; small seeds are eaten by birds; nectar is food for migrating monarch butterflies.
HABITAT: Ditches, savannas, marshes, wet meadows, pine barrens.
CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: High
pH: 4-7
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 504,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Heliopsis helianthoides
OXEYE SUNFLOWER

Native
Vigorous clump-forming species with a long bloom period; provides food and cover for birds.
HABITAT: Fields, woods, floodplains, streambanks.
CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 102,000
Seeding Rate: Up to 2% of a mix; of a meadow mix

Herbaceous Perennial



Hibiscus moscheutos
**CRIMSONEYED
ROSEMALLOW**

Native

Showy long-lived species in wet meadows and at the water's edge.

HABITAT: Alluvial meadows, swamp forest edges, brackish marshes.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

Pollinator Value: Low

pH: 4-7.5

Bloom Period: Summer/Fall

Flower Color(s): White

Approx Seeds Per Lb: 200,000

Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Hypericum pyramidatum
GREAT ST. JOHNSWORT

Native

Tall showy species with large flowers.

HABITAT: Alluvial shores, rocky banks, swamps.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

Pollinator Value: Low

pH: 5.7-7.1

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 1,800,000

Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Hypericum perforatum
**COMMON
ST. JOHNSWORT**

Naturalized

Source of herbal St. Johnswort; not for use in native mixes; seeds are persistent in soils.

HABITAT: Fields, roadsides, open spaces; tolerates poor soils.

CHARACTERISTICS:

Height: Up to 2.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 4,540,000

Seeding Rate: Up to 0.1% of a mix

Herbaceous Perennial



Kosteletzkya virginica
**VIRGINIA SALTMARSH
MALLOW**

Native

Attractive in coastal marshes.

HABITAT: Salt or brackish marshes, shores.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Summer/Fall

Flower Color(s): Pink

Approx Seeds Per Lb: 26,000

Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Hypericum punctatum
**SPOTTED
ST. JOHNSWORT**

Native

Commonly found in riparian areas.

HABITAT: Moist fields, floodplains, thickets, roadsides.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Shade

Drought Tolerance: Low

Pollinator Value: Low

pH: 4.6-7

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 2,000,000

Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Lathyrus sylvestris
FLAT PEA

Naturalized

Hardy rhizomatous legume; excellent for soil stabilization in infertile soils; seed must be incorporated into the soil to achieve successful establishment.

HABITAT: Borders of fields and thickets.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Shade

Drought Tolerance: High

pH: 5-7.8

Bloom Period: Summer

Flower Color(s): Purple, Pink, White

Approx Seeds Per Lb: 8,000

Seeding Rate: Up to 60% of a mix; up to 20 lb per acre alone with 15 lb per acre of tall fescue

Herbaceous Perennial Legume



Herbaceous Perennial Legume

Lespedeza capitata
ROUNDHEAD LESPEDEZA

Native
Provides food for birds and small ground animals.

HABITAT: Dry open woods, sand dunes, prairies.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 18 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
Pollinator Value: Low
pH: 5.7-8.2
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 174,000
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial

Linum perenne
PERENNIAL BLUE FLAX

Naturalized
Short stature and intense blue flowers make this a great species.

HABITAT: Naturalized along roadsides.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer
Flower Color(s): Blue

Approx Seeds Per Lb: 295,000
Seeding Rate: Up to 3% of a mix



Herbaceous Perennial Legume

Lespedeza virginica
SLENDER LESPEDEZA

Native
Clump-forming legume; our prettiest native Lespedeza; attractive flowers and foliage add color to native meadows; prolific producer of seeds eaten by game birds.

HABITAT: Open woods, roadsides, fields.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Low
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 175,000
Seeding Rate: Up to 3% of a mix; up to 10 PLS lb per acre alone



Herbaceous Perennial

Lobelia siphilitica
GREAT BLUE LOBELIA

Native
Decorative species with indeterminate blooms.

HABITAT: Swamps, moist meadows, streambanks, ditches.

CHARACTERISTICS:
Height: Up to 5.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Blue

Approx Seeds Per Lb: 7,760,000
Seeding Rate: Up to 0.3% of a mix



Herbaceous Perennial

Liatris spicata
MARSH BLAZING STAR

Native
Corm-forming species; flowers have a feathery appearance.

HABITAT: Moist fields, fencerows, roadsides.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: High
pH: 5.6-7.5
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 100,000
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial Legume

Lotus corniculatus
BIRD'S FOOT TREFOIL

Naturalized
Excellent for controlling erosion on strip mines and landfills; provides good livestock forage.

HABITAT: Fields, roadsides, meadows.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5-8
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 369,800
Seeding Rate: Up to 30% of a mix; up to 15 lb per acre alone



Ludwigia alternifolia
SEEDBOX

Native
Four-sided seedheads add texture to winter landscapes.

HABITAT: Wet woods, swampy fields.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,800,000
Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Lupinus polyphyllus
BIGLEAF LUPINE

Native
Very attractive in early summer; not for use in habitat restoration in our region.

HABITAT: Fields, roadsides; adapted to fine to coarse soils.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Bloom Period: Spring/Summer
Flower Color(s): Blue

Approx Seeds Per Lb: 75,000
Seeding Rate: Up to 2.5% of a mix

Herbaceous Perennial Legume



Ludwigia linearis
NARROWLEAF PRIMROSE WILLOW

Native
Fast-growing species with indeterminate blooms.

HABITAT: Savannas, ditches, bogs.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,000,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Lycopus americanus
AMERICAN WATER HOREHOUND

Native
Adds diversity to wet meadows and wetlands.

HABITAT: Wet ditches, swamps, moist thickets, fields, shaded hillsides.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Shade
Drought Tolerance: Low
Pollinator Value: High
pH: 5.2-7.8
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 3,025,300
Seeding Rate: Up to 0.4% of a mix

Herbaceous Perennial



Ludwigia maritima
SEASIDE PRIMROSE WILLOW

Native
Nice addition to a southeast coastal plain wetland mix.

HABITAT: Savannas, ditches, low pinelands.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,000,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Mimulus ringens
SQUARE STEMMED MONKEYFLOWER

Native
Rhizomatous species; reseeds itself in open wet areas; provides cover for wildlife; indeterminate snapdragon-like blooms are attractive to native pollinators.

HABITAT: Wet open ground of swamps, meadows, shores.

CHARACTERISTICS:
Height: Up to 4.2 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 22,900,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Monarda fistulosa
WILD BERGAMOT

Native
Showy rhizomatous species; often used as an ornamental.

HABITAT: Fields, brushy thickets, prairies, roadsides.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Minimum Root Depth: 4 in.
Shade Tolerance: Moderate
Drought Tolerance: No
Pollinator Value: High
pH: 6-8
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 1,272,500
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Oenothera biennis
EVENING PRIMROSE

Native
Shows the second year after seeding; new blossoms occur every evening and early morning during the season; provides good wildlife food and habitat, especially for birds.

HABITAT: Fields, prairies, roadsides, waste areas.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: Medium
pH: 5-7
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 1,376,000
Seeding Rate: Up to 0.2% of a mix

Herbaceous Biennial



Monarda punctata
SPOTTED BEEBALM

Native
Species with a pleasant fragrance.

HABITAT: Dry sandy and Coastal Plain soils, sandy upland forests, forest edges, fields, roadsides.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Unknown
Drought Tolerance: High
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 1,472,000
Seeding Rate: Up to 1% of a mix

Herbaceous Annual/Biennial/Perennial



Oenothera fruticosa var. fruticosa
SUNDROPS

Native
Attractive flowers.

HABITAT: Fields, meadows, roadsides, open woods.

CHARACTERISTICS:
Height: Up to 3.3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.5-7
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 3,780,000
Seeding Rate: Up to 0.6% of a mix

Herbaceous Perennial

DO THE MATH | *Calculate twice, order once.*

Ernst Seeds conditions and prepares seed to the highest standard and sells it in bulk or pure live seed (PLS) quantities. PLS refers to the percentage of live seed by weight in a seed lot having the potential to develop into a seedling. Calculate how much bulk seed you need by using the following calculations:

HOW TO CALCULATE PLS

Percent Total Germination =
(Germination + Hard Seed + Dormant Seed)

70 + 15 + 5 = 90% Total Germination

Use that figure and the purity percentage to calculate the PLS percentage.

Multiply Total Germination by Purity / 100 = PLS %
(95% X 90%)/100 = 85.5% of Pure Live Seed

What Does This Mean?

85.5% of this seed lot by weight has the potential to grow. If 10 pounds of pure live seed is needed on a site, the quantity of bulk seed needed is determined by:

100 / PLS = Pounds of bulk seed needed to produce 1 lb of PLS

100 / 85.5 = 1.16 lbs of bulk seed X 10 lbs = 11.6 lbs of bulk seed



Oenothera speciosa
**SHOWY EVENING
PRIMROSE**

Native

Large flowers open in the evening.

HABITAT: Dry open places, roadsides.

CHARACTERISTICS:

Height: Up to 2.6 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Spring/Summer

Flower Color(s): White, Pink

Approx Seeds Per Lb: 3,436,000

Seeding Rate: Up to 0.6% of a mix

Herbaceous Perennial



Penstemon hirsutus
HAIRY BEARDTONGUE

Native

Our shortest and earliest blooming Penstemon; attractive addition to short grass meadows.

HABITAT: Dry fields, woods, roadside banks, rocky slopes.

CHARACTERISTICS:

Height: Up to 3.3 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: High

Bloom Period: Spring/Summer

Flower Color(s): Purple

Approx Seeds Per Lb: 3,877,000

Seeding Rate: Up to 0.4% of a mix

Herbaceous Perennial



Papaver rhoeas
CORN POPPY

Naturalized

Creates an early field of color a year after seeding.

HABITAT: Roadsides, meadows, gardens.

CHARACTERISTICS:

Height: Up to 3 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Spring/Fall

Flower Color(s): Red

Approx Seeds Per Lb: 3,179,000

Seeding Rate: Up to 0.8% of a mix

Herbaceous Annual



Penstemon laevigatus
**APPALACHIAN
BEARDTONGUE**

Native

Provides early season bloom to meadows in full sun to partial shade.

HABITAT: Meadows, woods, roadsides.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: High

Bloom Period: Spring/Summer

Flower Color(s): Purple

Approx Seeds Per Lb: 350,000

Seeding Rate: Up to 4.5% of a mix

Herbaceous Perennial



Penstemon digitalis
**TALL WHITE
BEARDTONGUE**

Native

Durable early clump-forming species; found in many of our meadow mixes.

HABITAT: Meadows, old fields, roadsides.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 8 in.

Shade Tolerance: Shade

Drought Tolerance: High

Pollinator Value: High

pH: 5.5-7

Bloom Period: Spring/Summer

Flower Color(s): White

Approx Seeds Per Lb: 400,000

Seeding Rate: Up to 4% of a mix

Herbaceous Perennial



Penthorum sedoides
DITCH STONECROP

Native

Stoloniferous species; adds texture to wet landscapes; provides erosion control and habitat for wildlife.

HABITAT: Low wet ground, ditches.

CHARACTERISTICS:

Height: Up to 2.6 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

Pollinator Value: High

pH: 5-7

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 45,000,000

Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Herbaceous Annual

Polygonum pensylvanicum
PENNSYLVANIA SMARTWEED

Native
Acts as a cover crop in floodplain areas and on wetland sites; provides food and cover for wildlife.

HABITAT: Meadows, fields, waste places, moist ditches.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
Pollinator Value: Medium
pH: 4-8.5
Bloom Period: Spring/Fall
Flower Color(s): Pink

Approx Seeds Per Lb: 126,100
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Pycnanthemum tenuifolium
NARROWLEAF MOUNTAINMINT

Native
Rhizomatous mint species.

HABITAT: Dry soils of prairies and upland woods, moist old fields, floodplains, sandy streambanks.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer
Flower Color(s): White

Approx Seeds Per Lb: 5,336,000
Seeding Rate: Up to 0.4% of a mix



Herbaceous Annual/Perennial Vine

Polygonum sagittatum
ARROWLEAF TEARTHUMB

Native
Fast-growing vine; not for use in residential settings as the prickly stems can cut the skin; provides food for waterfowl.

HABITAT: Wet meadows, bogs, marshes.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
Pollinator Value: Medium
pH: 4-8.5
Bloom Period: Spring
Flower Color(s): Pink

Approx Seeds Per Lb: 125,000
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Pycnanthemum virginianum
VIRGINIA MOUNTAINMINT

Native
Rhizomatous mint species; stems and leaves are fragrant when crushed; also referred to as mountain thyme.

HABITAT: Upland woods, moist prairies.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 3,872,000
Seeding Rate: Up to 0.4% of a mix



Herbaceous Perennial

Pycnanthemum incanum
HOARY MOUNTAINMINT

Native
Rhizomatous, sweet-scented upland mint species; attractive to many diverse pollinators.

HABITAT: Upland woods, old fields, thickets, barrens.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): Purple

Seeding Rate: Up to 0.4% of a mix



Herbaceous Perennial

Ratibida columnifera
YELLOW PRAIRIE CONEFLOWER

Native
Attractive tap-rooted species in a perennial wildflower mix; source of food for birds and pollinators.

HABITAT: Dry open places, including prairies.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 5.9-7
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 737,100
Seeding Rate: Up to 2% of a mix



Ratibida pinnata
**GREY HEADED
CONEFLOWER**

Native

Long-lasting showy flowers; provides food and cover for wildlife.

HABITAT: Dry prairies, dry woods, old fields.

CHARACTERISTICS:

Height: Up to 4.8 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: Medium

pH: 5.6-6.8

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 427,500

Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Rudbeckia subtomentosa
**SWEET BLACKEYED
SUSAN**

Native

Provides food and cover for birds; nectar source for bees.

HABITAT: Prairies, low ground.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 712,000

Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Rudbeckia fulgida var. *fulgida*
ORANGE CONEFLOWER

Native

Provides decorative late summer to fall color.

HABITAT: Dry to moist woodlands, meadows.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Medium

Bloom Period: Summer/Fall

Flower Color(s): Yellow, Orange

Approx Seeds Per Lb: 500,000

Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Rudbeckia triloba
BROWNEYED SUSAN

Native

Provides late summer color and texture to landscapes; provides food for birds.

HABITAT: Old fields, rocky slopes, woodland edges.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Medium

Bloom Period: Summer/Fall

Flower Color(s): Yellow, Orange

Approx Seeds Per Lb: 536,000

Seeding Rate: Up to 0.5% of a mix

Herbaceous Biennial



Rudbeckia hirta
BLACKEYED SUSAN

Native

Most common native flower in our meadows; provides food and cover for birds.

HABITAT: Fields, meadows, roadsides.

CHARACTERISTICS:

Height: Up to 3.3 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: Medium

pH: 6-7

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 1,575,700

Seeding Rate: Up to 3% of a mix

Herbaceous Annual/Biennial/Perennial



Saururus cernuus
LIZARD'S TAIL

Native

Beautiful wetland wildflower.

HABITAT: Marshes, swamps.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Medium

Bloom Period: Summer

Flower Color(s): White

Approx Seeds Per Lb: 100,000

Seeding Rate: Up to 3% of a mix

Herbaceous Perennial



Herbaceous Perennial Legume

Senna hebecarpa
WILD SENNA

Native
Robust legume with sturdy growth; provides food for birds; excellent food source for bumblebees.

HABITAT: Streambanks, moist old fields, moist open woods.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,000
Seeding Rate: Up to 1.5% of a mix



Herbaceous Perennial

Silphium perfoliatum
CUP PLANT

Native
Robust long-lived legume; best established by planting in the fall as a dormant seeding with germination occurring in the spring; may be used as forage for domestic animals with multiple cuts; high potential as a bioenergy crop; provides food for birds and wildlife.

HABITAT: Floodplains, abandoned fields, moist meadows.

CHARACTERISTICS:
Height: Up to 9.8 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: High
pH: 4.5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 100,000
Seeding Rate: Up to 1% of a mix



Herbaceous Perennial Legume

Senna marilandica
MARYLAND SENNA

Native
Robust long-lived legume with sturdy growth; provides food and cover for wildlife.

HABITAT: Dry roadsides, rocky woodlands, streambanks.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: Medium
pH: 4-7
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,500
Seeding Rate: Up to 1.5% of a mix



Herbaceous Perennial

Silphium terebinthinaceum
PRAIRIE DOCK

Native
Adds texture to landscapes; seeds provide food for birds.

HABITAT: Prairies.

CHARACTERISTICS:
Height: Up to 13 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 17,000
Seeding Rate: Up to 5% of a mix



Herbaceous Perennial

Silphium asteriscus var. *laevicaule*
STARRY ROSINWEED

Native
Clump-forming legume; seeds provide food for birds.

HABITAT: Woodlands, old fields, thickets.

CHARACTERISTICS:
Height: Up to 8.8 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 20
Seeding Rate: Up to 2% of a mix



Herbaceous Perennial

Silphium trifoliatum
WHORLED ROSINWEED

Native
Clump-forming species; seeds provide food for birds; source of nectar for butterflies.

HABITAT: Roadsides, dry fields, meadows.

CHARACTERISTICS:
Height: Up to 9.8 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 20,800
Seeding Rate: Up to 1% of a mix



Solidago bicolor
WHITE GOLDENROD

Native

Clump-forming species; survives on low fertility sites and road cuts.

HABITAT: Dry woods, wooded banks, shale barrens.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Summer/Fall

Flower Color(s): White

Approx Seeds Per Lb: 1,649,000

Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Solidago nemoralis
GRAY GOLDENROD

Native

Our shortest goldenrod; food source for pollinators.

HABITAT: Fields, open woods, roadsides in low fertility soils.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: Very High

pH: 6.5-7.5

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 1,008,000

Seeding Rate: Up to 0.7% of a mix

Herbaceous Perennial



Solidago canadensis
CANADA GOLDENROD

Native

Aggressive rhizomatous species; does well with switchgrass and big bluestem; dominant robust vegetation adds diversity to native landscapes; provides cover for wildlife. (Note: Our harvest of this species from natural stands includes the species *Solidago gigantea* and *Solidago altissima*).

HABITAT: Moist or dry open places, fields, roadsides.

CHARACTERISTICS:

Height: Up to 8.1 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: Very High

pH: 4.8-7.5

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 4,600,000

Seeding Rate: Up to 0.2% of a mix

Herbaceous Perennial



Solidago odora
LICORICE SCENTED GOLDENROD

Native

Showy addition to landscapes; leaves have a scent resembling black licorice when crushed.

HABITAT: Dry open woods, sandy soils.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 2,268,000

Seeding Rate: Up to 0.3% of a mix

Herbaceous Perennial



Solidago juncea
EARLY GOLDENROD

Native

First goldenrod of the season to bloom; beautiful mid-summer addition to meadows.

HABITAT: Open woods, fields, meadows, roadsides.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Summer

Flower Color(s): Yellow

Approx Seeds Per Lb: 2,538,000

Seeding Rate: Up to 0.3% of a mix

Herbaceous Perennial



Solidago patula
ROUGHLEAF GOLDENROD

Native

Provides cover for wildlife.

HABITAT: Swamps, wet meadows, floodplains, moist woods.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 6 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

Pollinator Value: Very High

pH: 4.5-7

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 700,000

Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Solidago riddellii
RIDDELL'S GOLDENROD

Native
Attractive rhizomatous species; provides cover for wildlife.

HABITAT: Swamps, wet meadows, moist prairies.

CHARACTERISTICS:
Height: Up to 4.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 1,544,000
Seeding Rate: Up to 0.5% of a mix

Herbaceous Perennial



Solidago speciosa
SHOWY GOLDENROD

Native
Source of nectar for migrating monarch butterflies.

HABITAT: Open woods, prairies, fields, plains.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 1,340,000
Seeding Rate: Up to 0.6% of a mix

Herbaceous Perennial



Solidago rigida
STIFF GOLDENROD

Native
Attractive rhizomatous species with robust growth; provides food and cover for wildlife.

HABITAT: Dry open woods, prairies.

CHARACTERISTICS:
Height: Up to 6.5 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: High
Pollinator Value: Very High
pH: 5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 1,009,000
Seeding Rate: Up to 0.7% of a mix

Herbaceous Perennial



Verbena hastata
BLUE VERVAIN

Native
Easily seeded in wetlands.

HABITAT: Moist meadows, floodplains, wet ditches, roadsides.

CHARACTERISTICS:
Height: Up to 8.1 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Purple

Approx Seeds Per Lb: 1,488,000
Seeding Rate: Up to 4% of a mix

Herbaceous Biennial/Perennial



Solidago rugosa
WRINKLELEAF GOLDENROD

Native
Rhizomatous species; dense stems provide cover for wildlife.

HABITAT: Fields, woods, floodplains, thickets, roadsides, open ground.

CHARACTERISTICS:
Height: Up to 8.1 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: Very High
pH: 5-7.5
Bloom Period: Summer/Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 1,000,000
Seeding Rate: Up to 0.7% of a mix

Herbaceous Perennial



Verbena stricta
HOARY VERVAIN

Native
Adds texture and color to open landscapes.

HABITAT: Prairies, barrens, fields, roadsides; tolerates poor soils.

CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer
Flower Color(s): Purple

Approx Seeds Per Lb: 527,000
Seeding Rate: Up to 2% of a mix

Herbaceous Annual/Perennial



Verbena urticifolia
WHITE VERVAIN

Native
Adds texture to meadows.
HABITAT: Thickets, moist fields, meadows, open places.
CHARACTERISTICS:
Height: Up to 8.1 ft.
Shade Tolerance:
Drought Tolerance:
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): White
Approx Seeds Per Lb: 1,008,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Vernonia gigantea
GIANT IRONWEED

Native
Aggressive growth provides nesting habitat for woodcock; seeds are a source of food for birds.
HABITAT: Moist fields, wet woods, floodplains, meadows.
CHARACTERISTICS:
Height: Up to 9.8 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer/Fall
Flower Color(s): Purple
Approx Seeds Per Lb: 320,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Verbesina alternifolia
WINGSTEM

Native
Aggressive growth provides cover in wet meadows.
HABITAT: Moist wooded slopes, shaded lowlands, riverbanks.
CHARACTERISTICS:
Height: Up to 13 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer/Fall
Flower Color(s): Yellow
Approx Seeds Per Lb: 145,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Vernonia noveboracensis
NEW YORK IRONWEED

Native
Aggressive growth provides cover; seeds provide food for birds.
HABITAT: Streambanks, wet fields, pastures, meadows.
CHARACTERISTICS:
Height: Up to 8.1 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Moderate
Drought Tolerance: Moderate
Pollinator Value: High
pH: 4.5-8
Bloom Period: Summer/Fall
Flower Color(s): Purple
Approx Seeds Per Lb: 300,000
Seeding Rate: Up to 1% of a mix

Herbaceous Perennial



Vernonia angustifolia
TALL IRONWEED

Native
Showy clump-forming species.
HABITAT: Savannas, pine barrens, sandy woods, old fields.
CHARACTERISTICS:
Height: Up to 3.9 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: High
Bloom Period: Summer
Flower Color(s): Purple
Approx Seeds Per Lb: 300,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial



Veronicastrum virginicum
CULVER'S ROOT

Native
Showy rhizomatous species.
HABITAT: Moist meadows, thickets, swamps.
CHARACTERISTICS:
Height: Up to 6.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Pollinator Value: Very High
Bloom Period: Summer
Flower Color(s): Pink, White
Approx Seeds Per Lb: 7,761,000
Seeding Rate: Up to 0.1% of a mix

Herbaceous Perennial



Viola cornuta
JOHNNY JUMPUP

Naturalized
Pansy-like miniature species; added to mixes for early spring color.

HABITAT: Dry meadows.

CHARACTERISTICS:
Height: Up to 0.5 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Spring/Summer
Flower Color(s): White, Purple

Approx Seeds Per Lb: 796,000
Seeding Rate: Up to 2% of a mix

Herbaceous Annual



Zizia aurea
GOLDEN ALEXANDERS

Native
One of our earliest blooming natives; early source of food for pollinators.

HABITAT: Wooded bottomlands, streambanks, moist meadows, floodplains.

CHARACTERISTICS:
Height: Up to 3 ft.
Shade Tolerance: Moderate
Drought Tolerance: Unknown
Pollinator Value: Medium
Bloom Period: Spring/Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 172,000
Seeding Rate: Up to 2% of a mix

Herbaceous Perennial





2 mm Grid

Amorpha canescens
LEADPLANT

Native

Source of food for wildlife and pollinators.

HABITAT: Sandy open woods, dry prairies.

CHARACTERISTICS:

Height: Up to 3.9 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Shade

Drought Tolerance: High

pH: 5.5-8

Bloom Period: Summer

Flower Color(s): Purple

Approx Seeds Per Lb: 195,000

Seeding Rate: Up to 0.4% of a mix

Woody Deciduous Shrub



Woody Deciduous Shrub

Cephalanthus occidentalis
BUTTONBUSH

Native

Produces an abundance of seed favored by ducks; good source of nectar for butterflies and bees.

HABITAT: Low wet ground, swamps, bogs, streambanks, lake edges.

CHARACTERISTICS:

Height: Up to 9.8 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Shade

Drought Tolerance: Moderate

pH: 4.7-8.6

Bloom Period: Summer

Flower Color(s): White

Approx Seeds Per Lb: 134,000

Seeding Rate: Up to 0.6% of a mix; of a wetland mix



Woody Deciduous Shrub

Amorpha fruticosa
RIVER LOCUST

Native

Provides food and cover for wildlife; source of food for bees.

HABITAT: Alluvial soils along streams, rivers, other moist areas.

CHARACTERISTICS:

Height: Up to 16.3 ft.

Minimum Root Depth: 24 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

pH: 5-8.5

Bloom Period: Spring/Summer

Flower Color(s): Purple

Approx Seeds Per Lb: 77,000

Seeding Rate: Up to 1% of a mix



Woody Deciduous Shrub

Cornus amomum
SILKY DOGWOOD

Native

Thick, low vegetation provides excellent habitat for wildlife; abundant fruit is eaten by birds; blue berries in the fall.

HABITAT: Moist woods, fields, swamps, riparian areas.

CHARACTERISTICS:

Height: Up to 9.8 ft.

Minimum Root Depth: 16 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

Pollinator Value: Medium

pH: 5-7

Bloom Period: Spring/Summer

Flower Color(s): White

Approx Seeds Per Lb: 12,000

Seeding Rate: Up to 6.3% of a mix



Woody Deciduous Shrub

Amorpha herbacea
CLUSTERSPIKE FALSE INDIGO

Native

Attractive plant architecture and flowers.

HABITAT: Sandy fields, ridges, open woodlands; generally on the Coastal Plain.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Very High

Bloom Period: Spring/Summer

Flower Color(s): Blue, White

Approx Seeds Per Lb: 100,000

Seeding Rate: Up to 0.8% of a mix



Woody Deciduous Tree

Cornus florida
FLOWERING DOGWOOD

Native

Ornamental tree for woodland borders; provides abundant food for birds; red fruit.

HABITAT: Woods, woodland edges.

CHARACTERISTICS:

Height: Up to 39 ft.

Minimum Root Depth: 18 in.

Shade Tolerance: Shade

Drought Tolerance: Low

Pollinator Value: Medium

pH: 4.8-7.7

Bloom Period: Spring/Summer

Flower Color(s): Pink, White

Approx Seeds Per Lb: 4,500

Seeding Rate: Up to 5.6% of a mix



Woody Deciduous Shrub

Cornus racemosa
GRAY DOGWOOD

Native
Attractive rhizomatous shrub; thick, low vegetation provides excellent habitat for wildlife; fruit is a fall and winter food source for birds; white berries on red panicles.

HABITAT: Moist meadows, thickets, streambanks, roadsides.

CHARACTERISTICS:
Height: Up to 16.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Shade
Drought Tolerance: Moderate
Pollinator Value: High
pH: 4.8-7.4
Bloom Period: Summer
Flower Color(s): White

Approx Seeds Per Lb: 13,000
Seeding Rate: Up to 5.8% of a mix



Woody Deciduous Shrub

Ilex verticillata
WINTERBERRY

Native
Red berries provide fall and winter food for wildlife.

HABITAT: Swamps, bogs, moist woods, wet shores.

CHARACTERISTICS:
Height: Up to 16.3 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Pollinator Value: Low
pH: 4.5-7.5
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 92,000
Seeding Rate: Up to 0.8% of a mix



Woody Deciduous Tree/Shrub

Cornus sericea
RED OSIER DOGWOOD

Native
Attractive stoloniferous shrub; thick, low vegetation provides excellent habitat for wildlife; bright red stems add winter color; abundant white fruit is eaten by birds.

HABITAT: Moist woods, moist meadows, thickets, riparian areas, swamps.

CHARACTERISTICS:
Height: Up to 9.8 ft.
Minimum Root Depth: 20 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
Pollinator Value: Medium
pH: 4.8-7.5
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 18,000
Seeding Rate: Up to 4.2% of a mix



Woody Deciduous Vine

Parthenocissus quinquefolia
VIRGINIA CREEPER

Native
High-growing vine; produces black fruit in the fall.

HABITAT: Woods, fields, woodland edges.

CHARACTERISTICS:
Height: Up to 49 ft.
Minimum Root Depth: 16 in.
Shade Tolerance: Moderate
Drought Tolerance: High
Pollinator Value: Low
pH: 5-7.5
Bloom Period: Spring/Summer
Flower Color(s): Green

Approx Seeds Per Lb: 18,000
Seeding Rate: Up to 1% of a mix



Woody Deciduous Shrub

Hamamelis virginiana
WITCHHAZEL

Native
Provides mid-story habitat; nuts provide fall food for wildlife.

HABITAT: Moist woods, brushy fields.

CHARACTERISTICS:
Height: Up to 16.3 ft.
Minimum Root Depth: 20 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.5-6.2
Bloom Period: Fall
Flower Color(s): Yellow

Approx Seeds Per Lb: 11,000
Seeding Rate: Up to 6.8% of a mix



Woody Deciduous Tree

Platanus occidentalis
AMERICAN SYCAMORE

Native
Large riverbank tree; provides shade and cover for riparian shrubs and forbs.

HABITAT: Streambanks, low woods, floodplains.

CHARACTERISTICS:
Height: Up to 162.5 ft.
Minimum Root Depth: 30 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 4.9-6.5
Bloom Period: Spring/Summer

Approx Seeds Per Lb: 192,000
Seeding Rate: Up to 0.1% of a mix



Prunus serotina
WILD BLACK CHERRY

Native

One of North America's most valuable hardwoods; provides food (fruit) and nesting sites for wildlife; source of food for caterpillars of 300 species of moths and butterflies eaten by birds; purplish-black berries in the fall.

HABITAT: Woods, fencerows.

CHARACTERISTICS:

Height: Up to 97.5 ft.

Minimum Root Depth: 36 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: High

pH: 4-7.5

Bloom Period: Spring/Summer

Flower Color(s): White

Approx Seeds Per Lb: 4,800

Seeding Rate: Up to 5.2% of a mix

Woody Deciduous Tree



Robinia pseudoacacia
BLACK LOCUST

Native

Provides nitrogen and cover for successive vegetation; good honey producer.

HABITAT: Reclamation sites, floodplains, thickets, fencerows; develops quickly in poor soils.

CHARACTERISTICS:

Height: Up to 97.5 ft.

Minimum Root Depth: 36 in.

Shade Tolerance: Full Sun

Drought Tolerance: High

pH: 4.6-8.2

Bloom Period: Spring/Summer

Flower Color(s): White

Approx Seeds Per Lb: 24,000

Seeding Rate: Up to 1% of a mix

Woody Deciduous Tree



Prunus virginiana
CHOCHECHERRY

Native

Creates intermediate to mid-story habitat for nesting; abundant fruit provides summer and fall food for birds; host to many native caterpillars that are also food for birds; dark red to purple berries in the summer and fall.

HABITAT: Rocky upland woods, roadsides.

CHARACTERISTICS:

Height: Up to 32.5 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: High

pH: 5.2-8.4

Bloom Period: Spring/Summer

Flower Color(s): White

Approx Seeds Per Lb: 5,000

Seeding Rate: Up to 5% of a mix

Woody Deciduous Shrub



Rosa carolina
PASTURE ROSE

Native

Provides cover for wildlife.

HABITAT: Dunes, prairies, fields, upland woods.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Moderate

Drought Tolerance: High

Pollinator Value: Medium

pH: 4-7

Bloom Period: Spring/Summer

Flower Color(s): Pink

Approx Seeds Per Lb: 50,000

Seeding Rate: Up to 1.5% of a mix

Woody Deciduous Shrub



Rhus typhina
STAGHORN SUMAC

Native

Bark provides winter food for cottontail rabbits; red fruit is an emergency winter food source for birds, including wild turkey.

HABITAT: Dry open fields, roadsides, woodland edges.

CHARACTERISTICS:

Height: Up to 32.5 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Medium

Bloom Period: Summer

Flower Color(s): Green

Approx Seeds Per Lb: 60,000

Seeding Rate: Up to 1.3% of a mix

Woody Deciduous Shrub



Rosa palustris
SWAMP ROSE

Native

Attractive rhizomatous shrub; provides food and cover for songbirds, deer, ruffed grouse, pheasants, and wild turkey; develops rose hips that last into winter.

HABITAT: Streambanks, wetlands, marshes.

CHARACTERISTICS:

Height: Up to 8.1 ft.

Minimum Root Depth: 18 in.

Shade Tolerance: Shade

Drought Tolerance: Low

Pollinator Value: Medium

pH: 4-7

Bloom Period: Spring/Summer

Flower Color(s): Pink

Approx Seeds Per Lb: 24,970

Seeding Rate: Up to 3% of a mix

Woody Deciduous Shrub



Salix amygdaloides
PEACHLEAF WILLOW

Native
Good bioengineering material; fibrous roots grow into the water table to stabilize the streambank; leaf drops replenish organic life in streams; provides shade and habitat for wildlife.

HABITAT: Riparian areas, wetlands.

CHARACTERISTICS:

Height: Up to 65 ft.
Minimum Root Depth: 30 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 6-8
Bloom Period: Spring

Seeding Rate: Space live stakes on 3'-6' centers; may be used as larger live post plantings

Woody Deciduous Tree



Salix exigua ssp. interior
SANDBAR WILLOW

Native
Rhizomatous species; excellent bioengineering material; tolerant of ice and debris loading from streamflow; provides habitat for wildlife.

HABITAT: Sandbars, sandy or gravel streambanks, waterways.

CHARACTERISTICS:

Height: Up to 19.5 ft.
Minimum Root Depth: 36 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4-7.8
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Shrub



Salix discolor
PUSSY WILLOW

Native
Good bioengineering material; early show of catkins is the first sign of spring; provides habitat for wildlife.

HABITAT: Wetlands, moist or wet woods.

CHARACTERISTICS:

Height: Up to 22.8 ft.
Minimum Root Depth: 20 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4-7
Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Tree



Salix lucida
SHINING WILLOW

Native
Good bioengineering material; well-suited for wetland restoration; provides streambank erosion protection and habitat for wildlife.

HABITAT: Wetlands, streambanks.

CHARACTERISTICS:

Height: Up to 19.5 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5.8-7.2
Bloom Period: Spring
Flower Color(s): Yellow

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Shrub



Salix eriocephala
HEARTLEAF WILLOW

Native
Good rhizomatous bioengineering material; provides habitat for wildlife.

HABITAT: Riparian areas, streambanks.

CHARACTERISTICS:

Height: Up to 19.5 ft.
Minimum Root Depth: 20 in.
Shade Tolerance: Shade
Drought Tolerance: Low
pH: 4-7
Bloom Period: Spring/Summer

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Shrub



Salix nigra
BLACK WILLOW

Native
Stems can be brittle, making this only a fair bioengineering material; leaf drops replenish organic life in streams; provides shade and habitat for wildlife.

HABITAT: Wet meadows, riparian areas.

CHARACTERISTICS:

Height: Up to 65 ft.
Minimum Root Depth: 32 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4.8-8
Bloom Period: Spring/Summer

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Tree



Salix purpurea
STREAMCO WILLOW

Naturalized

Excellent early rooting bioengineering material; leaf drops replenish organic life in streams; provides shade to small streams and habitat for wildlife.

HABITAT: Streambanks, riparian areas.

CHARACTERISTICS:

Height: Up to 16.3 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Full Sun

Drought Tolerance: No

pH: 5.5-7.5

Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Shrub



Sambucus canadensis
ELDERBERRY

Native

Purplish-black berries provide an excellent source of summer food for wildlife.

HABITAT: Woods, moist fields, streambanks, moist roadsides.

CHARACTERISTICS:

Height: Up to 13 ft.

Minimum Root Depth: 16 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

Pollinator Value: Medium

pH: 5-8.9

Bloom Period: Summer

Flower Color(s): White, Yellow

Approx Seeds Per Lb: 292,000

Seeding Rate: Up to 0.3% of a mix

Woody Deciduous Shrub



Salix sericea
SILKY WILLOW

Native

Excellent bioengineering material; provides dense habitat for wildlife.

HABITAT: Wetlands, streambanks, riparian areas.

CHARACTERISTICS:

Height: Up to 13 ft.

Minimum Root Depth: 18 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

pH: 5.2-7

Bloom Period: Spring

Seeding Rate: Space live stakes on 3' centers

Woody Deciduous Shrub



Viburnum dentatum
ARROWWOOD

Native

Provides late summer and fall food, cover, browse, and nesting sites for birds; bluish-black fruit.

HABITAT: Wet woods, swamps.

CHARACTERISTICS:

Height: Up to 16.3 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Pollinator Value: Medium

Bloom Period: Spring/Summer

Flower Color(s): Pink, White

Approx Seeds Per Lb: 20,000

Seeding Rate: Up to 3.8% of a mix; space rooted seedlings on 6' centers

Woody Deciduous Shrub



Salix x cottetii
DWARF WILLOW

Naturalized

Excellent bioengineering material; ideal for utility stream crossings; tolerant of ice and debris loading from streamflow; resilient species that will recover from vehicle disturbance; provides good habitat for wildlife.

HABITAT: Streambanks.

CHARACTERISTICS:

Height: Up to 8 ft.

Minimum Root Depth: 20 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

pH: 5.5-7.5

Bloom Period: Spring

Seeding Rate: Space live stakes on 2' centers

Woody Deciduous Shrub



Viburnum lentago
NANNYBERRY

Native

Dense foliage provides cover; bluish-black fruit is a food source for wildlife.

HABITAT: Woods, wetlands, roadsides.

CHARACTERISTICS:

Height: Up to 32.5 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Shade

Drought Tolerance: Low

Pollinator Value: Medium

pH: 5-7

Bloom Period: Spring/Summer

Flower Color(s): Pink, White

Approx Seeds Per Lb: 7,843

Seeding Rate: Up to 9.6% of a mix

Woody Deciduous Shrub



Viburnum trilobum
AMERICAN CRANBERRY

Native
Red berries provide early winter food for wildlife.

HABITAT: Wetlands, wet woods.

CHARACTERISTICS:

Height: Up to 16.3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: No
Pollinator Value: Medium
pH: 5.5-7.5
Bloom Period: Spring/Summer
Flower Color(s): White

Approx Seeds Per Lb: 13,600
Seeding Rate: Up to 5.5% of a mix

Woody Deciduous Shrub





2 mm Grid

Herbaceous Perennial

Arrhenatherum elatius ssp. *elatius***TALL OATGRASS, 'RUFFNER'****Naturalized**

Bunchgrass with a pronounced cool season growth habit; used for forage and as a cover crop; good into Canada; highly palatable for whitetail deer throughout late fall and early winter.

HABITAT: Roadsides, fields, waste ground; persists in shallow, moderately infertile soils.

CHARACTERISTICS:

Height: Up to 6.5 ft.

Minimum Root Depth: 14 in.

Shade Tolerance: Moderate

Drought Tolerance: Moderate

pH: 5-7

Bloom Period: Summer

Approx Seeds Per Lb: 189,000

Seeding Rate: Up to 10 lb per acre when direct drilled; up to 20 lb per acre when broadcast seeded



Herbaceous Annual

Brassica napus
FORAGE RAPE, 'BONAR'**Naturalized**

Late maturing, high yielding forage rape; blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; quality winter feed for cattle and sheep; palatable for all livestock and grazing wildlife.

HABITAT: Pastures; grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 2.3 ft.

Minimum Root Depth: 6 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 6-7.2

Bloom Period: Spring/Summer

Seeding Rate: Up to 5 lb per acre when direct drilled



Herbaceous Annual

Avena sativa
OATS**Naturalized**

Small, bunch-type cereal grain for human and animal consumption; companion cover crop with a spring or fall seeding; matures quickly in hot weather and killed by freezing winter weather.

HABITAT: Tolerates a wide range of soil types, but prefers fertile open areas.

CHARACTERISTICS:

Height: Up to 4 ft.

Minimum Root Depth: 8 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 5.3-8.5

Bloom Period: Summer

Approx Seeds Per Lb: 19,000

Seeding Rate: Up to 90 lb per acre as a grain crop; up to 30 lb per acre as a cover crop with a perennial planting



2 mm Grid

Herbaceous Annual

Brassica napus
FORAGE RAPE, 'RANGIORA'**Naturalized**

Late maturing, high yielding forage rape; blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; quality winter feed for cattle and sheep; palatable for all livestock and grazing wildlife.

HABITAT: Pastures; grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 2.3 ft.

Minimum Root Depth: 6 in.


Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 6-7.2

Bloom Period: Spring/Summer

Seeding Rate: Up to 5 lb per acre when direct drilled



2 mm Grid

Herbaceous Annual/Biennial

Brassica napus
FORAGE BRASSICA, 'WINFRED'**Naturalized**

Blooms in the spring if planted in the fall; blooms 70 days post-planting if planted in the spring; leafy growth is palatable for deer, cattle, and sheep.

HABITAT: Food plots, forage (pastures)

CHARACTERISTICS:

Height: Up to 1.8 ft.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 5.5-7.5

Bloom Period: Spring

Flower Color(s): Yellow

Approx Seeds Per Lb: 157,000

Seeding Rate: Up to 8 lb per acre when direct drilled



Herbaceous Annual/Biennial

Brassica napus
WINTER CANOLA**Naturalized**

Can be very productive if planted in the fall; seeds are approximately 37% oil; also used in the production of healthy cooking oils; high quality livestock supplement; preferred by deer in the winter; makes good biodiesel.

HABITAT: Agricultural soils that support winter wheat production; grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 6 ft.

Minimum Root Depth: 6 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 6-7.2

Bloom Period: Spring

Flower Color(s): Yellow

Approx Seeds Per Lb: 160,000

Seeding Rate: Up to 10 lb per acre when direct drilled



Brassica napus
DWARF ESSEX RAPE

Naturalized

Member of the mustard family; seeds are approximately 37% oil; used in the production of healthy cooking oils; high quality livestock supplement; preferred by deer in the winter; makes good biodiesel.

HABITAT: Agricultural soils that support winter wheat production; grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 6 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
pH: 6-7.2
Bloom Period: Spring Yellow

Seeding Rate: Up to 10 lb per acre when direct drilled

Herbaceous Annual/Biennial



Brassica rapa
HYBRID FORAGE TURNIP, 'PACER'

Naturalized

Tap-rooted cross between forage rape and forage turnip; high energy, early maturing hybrid forage brassica; excellent regrowth (50-70 days); excellent for grazing.

HABITAT: Grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 2 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Bloom Period: Spring

Seeding Rate: Up to 5 lb per acre when direct drilled

Herbaceous Annual/Biennial



Brassica rapa
HYBRID FORAGE BRASSICA, 'VIVANT'

Naturalized

Tap-rooted brassica; used by dairy cattle, beef cattle, and sheep; graze within 35-50 days of planting and at 25-30-day intervals thereafter.

HABITAT: Grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 2 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Bloom Period: Spring

Seeding Rate: Up to 6 lb per acre when direct drilled

Herbaceous Annual/Biennial



Brassica rapa
PURPLE TOP TURNIP

Naturalized

Tap-rooted species; leaves are high in protein; highly palatable and digestible for deer.

HABITAT: Grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Bloom Period: Spring

Seeding Rate: Up to 4 lb per acre when direct drilled

Herbaceous Annual/Biennial



Brassica rapa
FORAGE TURNIP, 'APPIN'

Naturalized

Also referred to as field mustard; establishes in 50-80 days; 80% digestible and high in protein and carbohydrates; tolerates multiple grazings; multi-crowned for improved regrowth.


HABITAT: Pastures, fields, fertile open areas; grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
Bloom Period: Spring

Approx Seeds Per Lb: 192,800
Seeding Rate: Up to 5 lb per acre when direct drilled

Herbaceous Annual



Brassica spp.
BRASSICA, FORAGE, 'RANGI'

Naturalized

Provides food for wildlife.

HABITAT: Fields, waste ground.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5-8
Bloom Period: Spring

Seeding Rate: Up to 5 lb per acre when direct drilled

Herbaceous Annual/Biennial



Herbaceous Annual/Biennial

Camelina sativa CAMELINA

Naturalized

Member of the mustard family; used in the production of culinary oils and biodiesel; very low requirements for fertility, tillage, and weed control; oil and meal are high in Omega 3 fatty acids; approved by the USDA for human, poultry, and livestock consumption.

HABITAT: Prospers in various climates and soils; well-suited for marginal soils.

CHARACTERISTICS:

Height: Up to 3 ft.
Shade Tolerance: Unknown
Drought Tolerance: Unknown
Bloom Period: Summer
Flower Color(s): White, Yellow

Approx Seeds Per Lb: 400,000
Seeding Rate: Up to 6 lb per acre when direct drilled



Herbaceous Annual

Fagopyrum esculentum BUCKWHEAT

Naturalized

Tap-rooted, renovation grain cover crop for low productivity land; builds organic matter that decays rapidly when plowed under; planting time varies with application; killed by frost; provides food for human consumption, wildlife, and honey production.

HABITAT: Adapts to a wide range of soil types and conditions.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Moderate
Drought Tolerance: Low
pH: 5-8.5
Bloom Period: Summer

Approx Seeds Per Lb: 20,000
Seeding Rate: Up to 75 lb per acre alone (not recommended as a companion crop)



Herbaceous Perennial

Cichorium spp. CHICORY

Naturalized

High-yielding tap-rooted species; not for ornamental purposes; selected for intense grazing, less bolting, and better disease resistance; 20% protein; suitable for all wildlife and production livestock; excellent forage quality.

HABITAT: Grows best in moderately drained, medium to high fertility soils.

CHARACTERISTICS:

Height: Up to 2 ft.
Shade Tolerance: Unknown
Drought Tolerance: High
pH: 5.6-6.5
Bloom Period: Summer

Seeding Rate: Up to 2 lb per acre in a mix; up to 5 lb per acre alone



Herbaceous Perennial

Festuca elatior x Lolium perenne FESTULOLIUM, 'DUO'

Naturalized

Endophyte-free, high sugar cross between meadow fescue and a tetraploid perennial ryegrass; excellent palatability.

HABITAT: Fertile soils.

CHARACTERISTICS:

Height: Up to 2.7 ft.
Shade Tolerance: Unknown
Drought Tolerance: Moderate
Bloom Period: Summer

Approx Seeds Per Lb: 227,000
Seeding Rate: Up to 25 lb per acre in a mix; up to 30 lb per acre alone



Herbaceous Annual

Echinochloa crusgalli var. frumentacea JAPANESE MILLET

Naturalized

Warm season bunchgrass; used for erosion control and as a fast-growing summer companion crop; seed in the spring or summer; after growth, may be flooded to a depth of 18" during waterfowl migration season; provides food for wildlife and a favorite of waterfowl.

HABITAT: Well-drained soils, but thrives in wetlands.

CHARACTERISTICS:

Height: Up to 4.9 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 4.7-7.4
Bloom Period: Summer

Approx Seeds Per Lb: 143,000
Seeding Rate: Up to 30 lb per acre as a grain crop; up to 10 lb per acre as a companion crop



Herbaceous Annual Legume

Glycine max SOYBEAN, 'BOBWHITE'

Naturalized

Trailing legume for use in wildlife food plots; matures 120 days after planting.

HABITAT: Well to moderately drained soils and full sun.

CHARACTERISTICS:

Height: Up to 4 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 6.5-7.5
Bloom Period: Summer
Flower Color(s): Purple

Seeding Rate: Up to 60 lb per acre alone



Herbaceous Annual

Helianthus spp.
BLACK OIL SUNFLOWER

Naturalized

Grain is used for oil and meal; meal provides quality livestock feed; extremely popular food source for birds; may be left in the field for winter bird food.

HABITAT: Food plots.

CHARACTERISTICS:

Height: Up to 9.8 ft.
Minimum Root Depth: 8 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 0-6
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 7,000
Seeding Rate: Up to 10% of a food plot mix; up to 20 lb per acre alone or 0.5 lb per 1,000 sq ft



Herbaceous Biennial Legume

Melilotus officinalis
YELLOW BLOSSOM SWEETCLOVER

Naturalized

Tap-rooted legume; builds soil organic matter on low fertility sites; source of nectar and pollen for honeybees.

HABITAT: Mine spoil, roadsides, waste places.

CHARACTERISTICS:

Height: Up to 6.5 ft.
Minimum Root Depth: 32 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 5-8
Bloom Period: Summer
Flower Color(s): Yellow

Approx Seeds Per Lb: 259,000
Seeding Rate: Up to 5 lb per acre in a mix; up to 20 lb per acre alone



Herbaceous Perennial Legume

Medicago sativa
ALFALFA

Naturalized

Hardy component of right-of-way mixes; grows quickly after mowing; good livestock and wildlife forage.

HABITAT: Well-drained high fertility soils.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 24 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 6-8.5
Bloom Period: Spring
Flower Color(s): Purple

Approx Seeds Per Lb: 227,000
Seeding Rate: Up to 10 lb per acre in a mix; up to 20 lb per acre alone



Herbaceous Perennial Legume

Onobrychis viciifolia
SAINFOIN

Naturalized

Provides forage for deer, elk, cattle, and sheep; do not graze for two seasons following planting.

HABITAT: Hay fields, wildlife food plots; less persistent in moist soils.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 14 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 6-8.5
Bloom Period: Spring/Summer
Flower Color(s): Red

Approx Seeds Per Lb: 30,000
Seeding Rate: Up to 25% of a mix; up to 34 lb per acre alone



Herbaceous Biennial Legume

Melilotus alba
WHITE BLOSSOM SWEETCLOVER

Naturalized

Tap-rooted legume; source of nectar and pollen for bees.

HABITAT: Roadsides, waste places.

CHARACTERISTICS:

Height: Up to 9.8 ft.
Minimum Root Depth: 32 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 259,000
Seeding Rate: Up to 5 lb per acre in a mix; up to 20 lb per acre alone



Herbaceous Annual

Panicum miliaceum
WHITE PROSO MILLET

Naturalized

Grain for human and livestock consumption; seeds are one of the most attractive foods for birds.

HABITAT: Well-drained loamy soils; tolerates various soil conditions.

CHARACTERISTICS:

Height: Up to 3.3 ft.
Minimum Root Depth: 10 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.8-6.8
Bloom Period: Summer

Seeding Rate: Up to 35% of a mix; up to 20 lb per acre alone



Phleum pratense
TIMOTHY

Naturalized

Bunch-type grass; excellent forage for horses and other livestock.

HABITAT: Fields, open areas; moderate fertility requirements.

CHARACTERISTICS:

Height: Up to 4.9 ft.

Minimum Root Depth: 10 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

pH: 5.5-7

Bloom Period: Summer/Fall

Flower Color(s): Yellow

Approx Seeds Per Lb: 1,230,000

Seeding Rate: Up to 90% of a mix; up to 10 lb per acre alone

Herbaceous Perennial



Sanguisorba minor
SMALL BURNET

Naturalized

Very attractive to deer for its herbal taste.

HABITAT: Sunny flatlands to open slopes in well-drained soils.

CHARACTERISTICS:

Height: Up to 2.3 ft.

Minimum Root Depth: 12 in.

Shade Tolerance: Moderate

Drought Tolerance: Low

pH: 6-8

Bloom Period: Spring/Summer

Flower Color(s): Red

Approx Seeds Per Lb: 48,700

Seeding Rate: Up to 10% of a food plot mix; up to 2 lb per acre with clover

Herbaceous Perennial



Pisum arvense
AUSTRIAN WINTER PEA

Naturalized

Cold-tolerant cool season legume; used for erosion control as a cover crop or temporary hay crop; builds nitrogen and organic matter in fields and gardens; not winter hardy north of the Mason-Dixon Line; seed in early spring or fall; excellent for wild game food plots.

HABITAT: Prefers dry soils.

CHARACTERISTICS:

Height: Up to 4 ft.

Shade Tolerance: Unknown

Drought Tolerance: Unknown

Bloom Period: Spring

Seeding Rate: Up to 50% of a mix; up to 60 lb per acre alone

Herbaceous Annual Legume



Secale cereale
RYE

Naturalized

Bunch-type winter companion or cover crop; used for erosion control; may be planted anytime of year, but preferably in the fall or winter as rye has a strong ability to grow in cold weather.

HABITAT: More productive than other cereals in infertile sandy or acidic soils.

CHARACTERISTICS:

Height: Up to 5 ft.

Minimum Root Depth: 8 in.

Shade Tolerance: Full Sun

Drought Tolerance: Moderate

pH: 4.5-8.2

Bloom Period: Spring

Flower Color(s): Yellow

Approx Seeds Per Lb: 18,000

Seeding Rate: Up to 90 lb per acre as a grain crop; up to 30 lb per acre as a cover crop with a perennial planting

Herbaceous Annual



Raphanus sativus
RADISH, 'GROUNDHOG'

Naturalized

Species with a taproot that can reach a depth of 6'; root exudates help to suppress nematodes; plant at least 60 days prior to frost.

HABITAT: Cultivated; occasionally escapes to roadsides or old fields.

CHARACTERISTICS:

Height: Up to 3 ft.

Shade Tolerance: Full Sun

Drought Tolerance: Unknown

Bloom Period: Spring

Flower Color(s): Pink, Purple, White

Seeding Rate: Up to 10 lb per acre drilled with up to 30 lb per acre of grain rye

Herbaceous Annual



Setaria italica
GERMAN MILLET

Naturalized

Warm season bunchgrass; used as a cover crop and for pasture and haylage; good supplemental hay for cattle and sheep; seed in the spring or summer; provides food for wildlife.

HABITAT: Grows best in well-drained loamy soils with low moisture.

CHARACTERISTICS:

Height: Up to 5 ft.

Minimum Root Depth: 8 in.

Shade Tolerance: Full Sun

Drought Tolerance: Low

pH: 5.3-6.9

Bloom Period: Summer

Flower Color(s): White

Approx Seeds Per Lb: 217,000

Seeding Rate: Up to 30 lb per acre as a grain crop; up to 10 lb per acre as a cover crop

Herbaceous Annual



2 mm Grid

Herbaceous Annual

Sorghum spp. SORGHUM/RED MILO

Naturalized

Early maturing species; most attractive to mourning doves in the fall; makes emergency food for turkey, pheasants, and deer during the winter.

HABITAT: Cultivated fields.

CHARACTERISTICS:

Height: Up to 5 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: High
pH: 5.5-7.5
Bloom Period: Summer

Approx Seeds Per Lb: 27,000
Seeding Rate: Up to 50% of a mix; up to 30 lb per acre with a grain drill or when broadcast seeded



Herbaceous Annual Legume

Trifolium incarnatum CRIMSON CLOVER

Naturalized


Used as a winter companion or cover crop in pasture, hay, and silage mixes and for erosion control; seed in the fall or early spring south of I-64; plow under before the next crop is planted; provides food for honeybees.

HABITAT: Sandy and clay-like soils.

CHARACTERISTICS:

Height: Up to 2.6 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 6-7
Bloom Period: Spring/Summer
Flower Color(s): Red

Approx Seeds Per Lb: 150,000
Seeding Rate: Up to 10 lb per acre as a companion crop with a wildflower mix; up to 30 lb per acre alone



2 mm Grid

Herbaceous Annual Legume

Trifolium alexandrinum BERSEEM CLOVER

Naturalized

Forage legume for deer and livestock.

HABITAT: Hay fields, food plots.

CHARACTERISTICS:

Height: Up to 3 ft.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 6.5-8
Bloom Period: Summer
Flower Color(s): White

Approx Seeds Per Lb: 207,000
Seeding Rate: Up to 20% in a mix with alfalfa; up to 14 lb per acre drill seeded



Herbaceous Perennial Legume

Trifolium pratense RED CLOVER

Naturalized

Short-lived legume; used as hay, pasture, and silage for domestic animals, or as a cover crop that builds nitrogen and organic matter; flowers develop again after early summer cutting; may be frost seeded during early spring; provides food for bumblebees.

HABITAT: Medium fertility soils.

CHARACTERISTICS:

Height: Up to 2.6 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 6-7.6
Bloom Period: Spring/Summer
Flower Color(s): Pink

Approx Seeds Per Lb: 272,000
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre alone



Herbaceous Perennial Legume

Trifolium hybridum ALSIKE CLOVER

Naturalized

Winter-hardy legume; provides forage for domestic animals and food and cover for wildlife.

HABITAT: Meadows, disturbed areas; tolerates low fertility.

CHARACTERISTICS:

Height: Up to 4 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.6-7.5
Bloom Period: Spring/Summer
Flower Color(s): Pink, White

Approx Seeds Per Lb: 680,000
Seeding Rate: Up to 25% of a mix; up to 10 lb per acre alone



Herbaceous Perennial Legume

Trifolium repens WHITE CLOVER

Naturalized

Good, stoloniferous, erosion control cover crop; grows again after mowing or grazing; highly palatable as food for wildlife and domestic animals; source of food for honeybees.

HABITAT: Moist soils, lawns, field borders.

CHARACTERISTICS:

Height: Up to 1 ft.
Minimum Root Depth: 12 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.2-8
Bloom Period: Summer/Fall
Flower Color(s): White

Approx Seeds Per Lb: 712,000
Seeding Rate: Up to 10% of a mix; up to 10 lb per acre alone



2 mm Grid

Urochloa ramosa
BROWN TOP MILLET

Naturalized

Short-lived species for pasture, hay, or wildlife forage; establishes quickly for erosion control; plant from May to August; matures in 60 days.

HABITAT: Tolerates low fertility acidic soils.

CHARACTERISTICS:

Height: Up to 3 ft.
Minimum Root Depth: 24 in.
Shade Tolerance: Full Sun
Drought Tolerance: Low
pH: 5.5-6.9
Bloom Period: Summer

Approx Seeds Per Lb: 75,000
Seeding Rate: Up to 20 lb per acre when direct drilled; up to 30 lb per acre when broadcast seeded; up to 10 lb per acre as a cover crop

Herbaceous Annual



Vicia villosa
HAIRY VETCH

Naturalized

Builds nitrogen and organic matter in fields and gardens; seed in early fall after row crop is harvested; likely to freeze at -30 degrees F.

HABITAT: Well-drained soils.

CHARACTERISTICS:

Height: Up to 7 ft.
Minimum Root Depth: 6 in.
Shade Tolerance: Full Sun
Drought Tolerance: Moderate
pH: 6-7.5
Bloom Period: Spring
Flower Color(s): Purple

Approx Seeds Per Lb: 16,000
Seeding Rate: Up to 10 lb per acre in a mix with up to 50 lb per acre of grain rye

Herbaceous Biennial Legume



INDEX TO SCIENTIFIC/Common NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>Achillea millefolium</i>	94	<i>australis</i>	98	<i>albolutescens</i>	80
<i>Acorus americanus</i>	78	<i>pendula</i>	98	<i>comosa</i>	81
<i>Agastache foeniculum</i>	94	<i>tinctoria</i>	98	<i>crinita</i>	81
<i>Agrimonia parviflora</i>	94	Barnyardgrass	84	<i>frankii</i>	81
<i>Agrimony, Small Flowered</i>	94	Beardtongue		<i>grayi</i>	81
<i>Agrostis</i>		Appalachian	111	<i>intumescens</i>	81
<i>alba</i>	78	Hairy	111	<i>lupulina</i>	81
<i>hyemalis</i>	78	Tall White	111	<i>lurida</i>	82
<i>perennans</i>	78	Beebalm, Spotted	110	<i>pensylvanica</i>	82
<i>stolonifera</i>	78	Beeblossom, Biennial	105	<i>scoparia</i>	82
<i>tenuis</i>	78	Beggartick	99	<i>shortiana</i>	82
Alfalfa	128	Bentgrass		<i>stipata</i>	82
<i>Alisma subcordatum</i>	79	Autumn	78	<i>stricta</i>	82
Alkaligrass	89	Creeping	78	<i>vulpinoidea</i>	83
<i>Allium cernuum</i>	79	Colonial	78	<i>Centaurea cyanus</i>	99
<i>Alopecurus arundinaceus</i>	79	Winter	78	<i>Cephalanthus occidentalis</i>	119
<i>Amorpha</i>		Bergamot, Wild	110	<i>Chamaecrista</i>	
<i>canescens</i>	119	<i>Bidens</i>		<i>fasciculata</i>	99
<i>fruticosa</i>	119	<i>aristosa</i>	98	<i>nictitans</i>	99
<i>herbacea</i>	119	<i>cernua</i>	99	<i>Chasmanthium latifolium</i>	83
<i>Amsonia ciliata</i>	94	<i>frondosa</i>	99	<i>Cheiranthus allionii</i>	100
<i>Andropogon</i>		Black Cherry, Wild	121	Chicory	
<i>gerardii</i>	79	Blackeyed Susan	113	Blue	100
<i>ternarius</i>	79	Sweet	113	Chokecherry	121
<i>virginicus</i>	79	Blanketflower	105	<i>Chrysanthemum maximum</i>	100
<i>Anemone</i>		Blazing Star, Marsh	108	Chufa	83
<i>canadensis</i>	94	<i>Blephilia ciliata</i>	99	<i>Cichorium</i>	
Canadian	94	Blue Eyed Grass, Narrowleaf	91	<i>intybus</i>	100
<i>virginiana</i>	94	Blueflag	87	<i>spp.</i>	127
<i>Anthoxanthum odoratum</i>	80	Bluegrass		<i>Cinna arundinacea</i>	83
<i>Apocynum cannabinum</i>	95	Fowl	89	<i>Clematis virginiana</i>	100
<i>Aquilegia canadensis</i>	95	Rough	89	Clover	
<i>Arrhenatherum elatius ssp. elatius</i>	125	Bluejoint, Canada	80	Alsiike	130
Arrow Arum	89	Bluestar, Fringed	94	Berseem	130
Arrowwood	123	Bluestem		Crimson	130
<i>Asclepias</i>		Big	79	Red	130
<i>incarnata</i>	95	Little	90	White	130
<i>syriaca</i>	95	Splitbeard	79	Columbine, Eastern	95
<i>tuberosa</i>	95	Boneset	104	Coneflower	
<i>Aster</i>		Bottlebrush Grass	84	Grey Headed	113
Aromatic	97	<i>Bouteloua</i>		Orange	113
Arrowleaf	97	<i>curtipendula</i>	80	Pale Purple	103
Bigleaf	96	<i>gracilis</i>	80	Purple	103
Calico	96	<i>Brassica</i>		Yellow Prairie	112
<i>divaricatus</i>	95	Forage, 'Rangi'	126	Cordgrass, Prairie	92
Flat Topped White	98	Forage, 'Winfred'	125	<i>Coreopsis</i>	
Heath	97	Hybrid Forage, 'Vivant'	126	<i>basalis</i>	101
<i>laevis</i>	96	<i>napus</i>	125	<i>grandiflora</i>	101
Lance Leaved	96	<i>rapa</i>	126	Lanceleaf	101
<i>lanceolatus</i>	96	<i>spp.</i>	126	<i>lanceolata</i>	101
<i>lateriflorus</i>	96	Brome, Smooth	71	Plains	101
Lowrie's Blue Wood	96	<i>Bromus inermis</i>	71	Tall	101
<i>lowrieanus</i>	96	Broomsedge	79	<i>tinctoria</i>	101
<i>macrophyllus</i>	96	Brown-eyed Susan	113	<i>tripteris</i>	101
New England	96	Buckwheat	127	Cornflower	99
<i>novae-angliae</i>	96	Bulrush		<i>Cornus</i>	
<i>oblongifolius</i>	97	Brown	91	<i>amomum</i>	119
<i>pilosus</i>	97	Green	90	<i>florida</i>	119
<i>preanthoides</i>	97	Northern	90	<i>racemosa</i>	120
<i>puniceus</i>	97	Softstem	91	<i>sericea</i>	120
Purplestem	97	Wood	90	<i>Coronilla varia</i>	101
<i>sagittifolius</i>	97	Bundleflower, Illinois	103	Cosmos	102
Showy	97	Bur Marigold	98	<i>bipinnatus</i>	102
Smooth Blue	96	Nodding	98,99	Sulphur	102
<i>spectabilis</i>	97	Bur Reed		<i>sulphureus</i>	102
<i>umbellatus</i>	98	Eastern	91	Cranberry, American	124
White Wood	95	Giant	91	Crownvetch	101
Zigzag	97	Burnet, Small	129	Cup Plant	114
<i>Avena sativa</i>	125	Buttonbush	119	Cutgrass, Rice	87
Avens, White	105	<i>Calamagrostis canadensis</i>	80	Culver's Root	117
Baby's Breath, Annual	105	Camelina	127	<i>Cyperus esculentus</i>	83
Bachelor's Button	99	<i>Camelina sativa</i>	127	<i>Dactylis glomerata</i>	84
<i>Baptisia</i>		Canola, Winter	125	Daisy, Shasta	100
<i>alba</i>	98	<i>Carex</i>		<i>Dalea</i>	

INDEX TO SCIENTIFIC/Common NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>candida</i>	102	Annual	105	Slender	108
<i>purpurea</i>	102	<i>aristata</i>	105	<i>virginica</i>	108
<i>Daucus carota</i>	102	Perennial	105	<i>Liatris spicata</i>	108
Deertongue	88	<i>pulchella</i>	105	<i>Linum perenne</i>	108
<i>Delphinium ajacis</i>	102	Gamagrass, Eastern	93	Lizard's Tail	113
<i>Deschampsia cespitosa</i>	84	<i>Gaura biennis</i>	105	Lobelia	
<i>Desmanthus illinoensis</i>	102	<i>Geum canadense</i>	105	Great Blue	108
<i>Desmodium</i>		<i>Glyceria</i>		<i>siphilitica</i>	108
<i>canadense</i>	103	<i>grandis</i>	86	Locust	
<i>paniculatum</i>	103	<i>striata</i>	86	Black	120
<i>Dianthus barbatus</i>	103	<i>Glycine max</i>	127	River	119
Dock, Prairie	114	Golden Alexanders	118	<i>Lolium multiflorum</i>	87
Dogwood		Goldenrod		<i>Lotus corniculatus</i>	108
Flowering	119	Canada	115	Lovegrass	
Gray	120	Early	115	Bigtop	85
Red Osier	120	Grassleaf	105	Purple	85
Silky	119	Gray	115	Weeping	85
Dropseed		Licorice Scented	115	<i>Ludwigia</i>	
Prairie	92	Riddell's	116	<i>alternifolia</i>	109
Rough	92	Roughleaf	115	<i>linearis</i>	109
Sand	92	Showy	116	<i>maritima</i>	109
Duck Potato	90	Stiff	116	Lupine, Bigleaf	109
<i>Echinacea</i>		White	115	<i>Lupinus polyphyllus</i>	109
<i>pallida</i>	103	Wrinkleleaf	116	<i>Lycopus americanus</i>	109
<i>purpurea</i>	103	Gramma		Mannagrass	
Echinochloa		Blue	80	American	86
<i>crusgalli</i> var. <i>frumentacea</i>	127	Sideoats	80	Fowl	86
<i>muricata</i>	84	<i>Gypsophila elegans</i>	105	<i>Medicago sativa</i>	128
Elderberry	123	Hairgrass, Tufted	84	<i>Melilotus</i>	
<i>Elymus</i>		<i>Hamamelis virginiana</i>	120	<i>alba</i>	128
<i>canadensis</i>	84	<i>Helenium</i>		<i>officinale</i>	128
<i>hystrix</i>	84	<i>autumnale</i>	106	Milkweed	
<i>riparius</i>	84	<i>flexuosum</i>	106	Butterfly	95
<i>virginicus</i>	85	<i>Helianthus</i>		Common	95
<i>Eragrostis</i>		<i>angustifolius</i>	106	Swamp	95
<i>curvula</i>	85	<i>annuus</i>	106	Millet	
<i>hirsuta</i>	85	<i>maximilianii</i>	106	Brown Top	131
<i>spectabilis</i>	85	spp.	128	German	129
<i>Eryngium yuccifolium</i>	104	<i>Heliopsis helianthoides</i>	106	Japanese	127
<i>Eupatorium</i>		<i>Hibiscus moscheutos</i>	107	White Proso	128
<i>coelestinum</i>	104	Horehound, American Water	109	Milo, Red	109
<i>fistulosum</i>	104	<i>Hypericum</i>		<i>Mimulus ringens</i>	104
<i>maculatum</i>	104	<i>perforatum</i>	107	Mistflower	106
<i>perfoliatum</i>	104	<i>punctatum</i>	107	<i>Monarda</i>	
<i>purpureum</i>	104	<i>pyramidatum</i>	107	<i>fistulosa</i>	110
<i>Euthamia graminifolia</i>	105	Hyssop		<i>punctata</i>	110
Evening Primrose	110	Anise	94	Monkeyflower, Square Stemmed	109
Showy	111	Lavender	94	Mountainmint	
<i>Fagopyrum esculentum</i>	128	<i>Ilex verticillata</i>	120	Hoary	112
False Indigo		Indian Blanket	105	Narrowleaf	112
Blue	98	Indiangrass	91	Virginia	112
Clusterspike	119	Indianhemp	95	Nannyberry	123
Yellow	98	<i>Iris versicolor</i>	87	Oatgrass, Tall, 'Ruffner'	125
Fern, Sensitive	88	Ironweed		Oats	
Fescue		Giant	117	River	83
Blue	86	New York	117	<i>Oenothera</i>	
Chewings	86	Tall	117	<i>biennis</i>	110
Creeping Red	86	Joe Pye Weed	104	<i>fruticosa</i> var. <i>fruticosa</i>	110
Hard	86	Purple Node	104	<i>speciosa</i>	111
Sheep	85	Spotted	104	Onion, Nodding	79
Tall	85	Johnny Jumpup	118	<i>Onobrychis viciifolia</i>	128
<i>Festuca</i>		<i>Juncus</i>		<i>Onoclea sensibilis</i>	88
<i>arundinacea</i>	85	<i>effusus</i>	87	Orchardgrass	84
<i>elatior</i> x <i>Lolium perenne</i>	85	<i>tenuis</i>	87	Pagoda Plant, Downy	99
<i>ovina</i>	85	Junegrass	87	Panicgrass	
<i>ovina</i> var. <i>duriuscula</i>	86	<i>Koeleria macrantha</i>	87	Beaked	88
<i>ovina</i> var. <i>glauca</i>	86	<i>Kosteletzkya virginica</i>	107	Coastal	88
<i>rubra</i>	86	Larkspur, Rocket	102	Redtop	88
<i>rubra</i> ssp. <i>commutata</i>	86	<i>Lathyrus sylvestris</i>	107	Smooth	88
Festulolum, 'Duo'	127	Leadplant	119	<i>Panicum</i>	
Flat Pea	107	<i>Leersia oryzoides</i>	87	<i>amarum</i>	88
Flax, Perennial Blue	108	<i>Lespedeza</i>		<i>anceps</i>	88
Foxtail, Creeping	79	<i>capitata</i>	108	<i>clandestinum</i>	88
<i>Gaillardia</i>		Roundhead	108	<i>dichotomiflorum</i>	88

INDEX TO SCIENTIFIC/Common NAMES

NAME	PAGE	NAME	PAGE	NAME	PAGE
<i>miliaceum</i>	128	Path	87	spp.	130
<i>rigidulum</i>	88	Soft	87	Soybean, 'Bobwhite'	127
<i>virgatum</i>	89	Rye	129	<i>Sparganium</i>	
<i>Papaver rhoeas</i>	111	Ryegrass, Annual	87	<i>americanum</i>	91
<i>Parthenocissus quinquefolia</i>	120	<i>Sagittaria latifolia</i>	90	<i>eurycarpum</i>	91
Pea		Sainfoin	128	<i>Spartina pectinata</i>	92
Austrian Winter	129	<i>Salix</i>		Spiderwort	
Partridge	99	<i>amygdaloides</i>	122	Ohio	92
Sensitive	99	<i>discolor</i>	122	Virginia	93
<i>Peltandra virginica</i>	89	<i>eriocephala</i>	122	Zigzag	92
<i>Penstemon</i>		<i>exigua ssp. interior</i>	122	<i>Sporobolus</i>	
<i>digitalis</i>	111	<i>lucida</i>	122	<i>asper</i>	92
<i>hirsutus</i>	111	<i>nigra</i>	122	<i>cryptandrus</i>	92
<i>laevigatus</i>	111	<i>purpurea</i>	123	<i>heterolepis</i>	92
<i>Penthorum sedoides</i>	111	<i>sericea</i>	123	St. Johnswort	
<i>Phleum pratense</i>	129	<i>x cottetii</i>	123	Common	107
Pickeralweed	89	Saltmarsh Mallow, Virginia	107	Great	107
<i>Pisum arvense</i>	129	<i>Sambucus canadensis</i>	123	Spotted	107
Plantain, Mud	79	<i>Sanguisorba minor</i>	129	Stonecrop, Ditch	111
<i>Platanus occidentalis</i>	120	<i>Saururus cernuus</i>	113	Sumac, Staghorn	121
<i>Poa</i>		<i>Schizachyrium scoparium</i>	90	Sundrops	110
<i>palustris</i>	89	<i>Scirpus</i>		Sunflower	
<i>trivialis</i>	89	<i>atrovirens</i>	90	Black Oil	128
<i>Polygonum</i>		<i>cyperinus</i>	90	Common	106
<i>pensylvanicum</i>	112	<i>expansus</i>	90	Maximilian's	106
<i>sagittatum</i>	112	<i>hattorianus</i>	90	Narrowleaf	106
<i>Pontederia cordata</i>	89	<i>pendulus</i>	91	Oxeye	106
Poppy, Corn	111	<i>validus</i>	91	Sweetclover	
Prairie Clover		<i>Secale cereale</i>	129	White Blossom	128
Purple	102	Sedge		Yellow Blossom	128
White	102	Awl	82	Sweetflag	78
Primrose Willow		Blunt Broom	82	Sweetwilliam	103
Narrowleaf	109	Cosmos	81	Switchgrass	89
Seaside	109	Fox	83	Sycamore, American	120
<i>Prunus</i>		Frank's	81	Tearthumb, Arrowleaf	112
<i>serotina</i>	121	Fringed	81	Thimbleweed	94
<i>virginiana</i>	121	Gray's	81	Tickseed	
<i>Puccinellia distans</i>	89	Greenwhite	80	Goldenmane	101
Purpletop	93	Hop	81	Largeflower	101
<i>Pycnanthemum</i>		Lurid	81	Tickseed Sunflower, Showy	98
<i>Incanum</i>	112	Pennsylvania	82	Ticktrefoil	
<i>tenuifolium</i>	112	Short's	82	Panicleleaf	103
<i>virginianum</i>	112	Star	81	Showy	103
Queen Anne's Lace	102	Tussock	82	Timothy	129
Radish, 'GroundHog'	129	Seedbox	109	<i>Tradescantia</i>	
Rape		<i>Senna</i>		<i>ohiensis</i>	92
Dwarf Essex	126	<i>hebecarpa</i>	114	<i>subaspera</i>	92
Forage, 'Bonar'	125	<i>marilandica</i>	114	<i>virginiana</i>	93
Forage, 'Rangiora'	125	Maryland	114	Trefoil, Bird's Foot	108
<i>Raphanus sativus</i>	129	Wild	114	<i>Tridens flavus</i>	93
Ratibida		<i>Setaria italica</i>	129	<i>Trifolium</i>	
<i>columnifera</i>	112	<i>Silphium</i>		<i>alexandrinum</i>	130
<i>pinnata</i>	113	<i>asteriscus var. laevicaule</i>	114	<i>hybridum</i>	130
Rattlesnake Master	104	<i>perfoliatum</i>	114	<i>incarnatum</i>	130
Redtop	78	<i>terebinthaceum</i>	114	<i>pratense</i>	130
Reedgrass, Wood	83	<i>trifoliatum</i>	114	<i>repens</i>	130
<i>Rhus typhina</i>	121	<i>Sisyrinchium angustifolium</i>	91	<i>Tripsacum dactyloides</i>	93
<i>Robinia pseudoacacia</i>	121	Smartweed, Pennsylvania	112	Turnip	
<i>Rosa</i>		Sneezeweed		Forage, 'Appin'	126
<i>carolina</i>	121	Common	106	Hybrid Forage, 'Pacer'	126
<i>palustris</i>	121	Purplehead	106	Purple Top	126
Rose		<i>Solidago</i>		<i>Urochloa ramosa</i>	131
Pasture	121	<i>bicolor</i>	115	<i>Verbena</i>	
Swamp	121	<i>canadensis</i>	115	<i>hastata</i>	116
Rosemallow, Crimoneyed	107	<i>juncea</i>	115	<i>stricta</i>	116
Rosinweed		<i>nemoralis</i>	115	<i>urticifolia</i>	117
Starry	114	<i>odora</i>	115	<i>Verbesina alternifolia</i>	117
Whorled	114	<i>patula</i>	115	Vernalgrass, Sweet	80
<i>Rudbeckia</i>		<i>riddellii</i>	116	<i>Vernonia</i>	
<i>fulgida var. fulgida</i>	113	<i>rigida</i>	116	<i>angustifolia</i>	117
<i>hirta</i>	113	<i>rugosa</i>	116	<i>gigantea</i>	117
<i>subtomentosa</i>	113	<i>speciosa</i>	116	<i>noveboracensis</i>	117
<i>triloba</i>	113	<i>Sorghastrum nutans</i>	91	<i>Veronicastrum virginicum</i>	117
Rush		Sorghum		Vervain	

INDEX TO SCIENTIFIC/COMMON NAMES

NAME	PAGE
Blue	116
Hoary	116
White	117
Vetch, Hairy	131
<i>Viburnum</i>	
<i>dentatum</i>	123
<i>lentago</i>	123
<i>trilobum</i>	124
<i>Vicia villosa</i>	131
<i>Viola cornuta</i>	118
Virgin's Bower	100
Virginia Creeper	120
Wallflower	100
Wild Indigo	
Largeleaf	98
White	98
Wildrice	93
Wildrye	
Canada	84
Riverbank	84
Virginia	84
Willow	
Black	122
Dwarf	123
Heartleaf	122
Peachleaf	122
Pussy	122
Sandbar	122
Shining	122
Silky	123
Streamco	123
Wingstem	117
Winterberry	120
Witchhazel	120
Woolgrass	90
Yarrow, Common	94
<i>Zizania aquatica</i>	93
<i>Zizia aurea</i>	118



ACKNOWLEDGMENTS

SOME INFORMATION IN THIS CATALOG COURTESY OF:

- Bell, C. Ritchie and Taylor, Bryan J. Florida Wild Flowers and Roadside Plants. Laurel Hill Press, 1982.
- Brown, Melvin L. and Brown, Russell G. Herbaceous Plants of Maryland. Port City Press, Inc., 1984.
- Brown, Russell G. and Brown, Melvin L. Woody Plants of Maryland. Port City Press. 1972.
- Clewell, Andre F. Guide to the Vascular Plants of the Florida Panhandle. Florida State University Press, 1985.
- Duncan, Wilbur H. and Kartesz, John T. Vascular Flora of Georgia - An Annotated Checklist. University of Georgia Press, 1981.
- Flora of North America Editorial Committee. Flora of North America Association. Oxford University Press. 2006.
- Gleason Ph.D., Henry A. and Cronquist, Ph.D., Arthur. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. 2nd ed. New York Botanical Garden, 1991.
- Magee, Dennis W. and Ahles, Harry E. Flora of the Northeast: A Manual of the Vascular Flora of New England and Adjacent New York. University of Massachusetts Press. 1999.
- Mohlenbrock, R.H. Guide to the Vascular Flora of Illinois. Southern Illinois University Press. 1986.
- Radford, Albert E., Ahles, Harry E. and Bell, C. Ritchie. Manual of the Vascular Flora of the Carolinas. University of North Carolina Press, 1964.
- Rhoads, Ann. Fowler & Block, Timothy A. The Plants of Pennsylvania. University of Pennsylvania Press, 2000.
- Rhoads, Ann Fowler and Klein, Jr., William McKinley. The Vascular Flora of Pennsylvania: Annotated Checklist and Atlas. American Philosophical Society, 1993.
- Richardson, L.L.; Adler, L.S.; Andicochea, J.; Regan, K.H.; Anthony, W.; Manson, J.S.; and Irwin, R.E. Secondary metabolites in floral nectar reduce parasite infections in bumblebees. Proceedings of the Royal Society B. 282:20142471,2015.
- Taylor, Walter Kingsley. Florida Wildflowers in Their Natural Communities. University Press of Florida, 1998.
- Taylor, Walter Kingsley. The Guide to Florida Wildflowers. Taylor Trade Publishing, 1992.
- The Great Plains Flora Association. Flora of the Great Plains. University Press of Kansas. 1986.
- Tobe, Dr. John D.; Burks, Kathy Craddock; Cantrell, Richard W.; Garland, Mark A.; Sweeley, Maynard E.; Hall, Dr. David W.; Wallace, Pete; Anglin, Guy; Nelson, Gil; Cooper, Dr. James R.; Bickner, David; Gilbert, Katherine, Aymond, Neil; Greenwood, Ken; and Raymond, Nina. Florida Wetland Plants - An Identification Manual. University of Florida, 1998.
- USDA, NRCS (2015). The PLANTS Database (<http://plants.usda.gov>, 23 October 2015). National Plant Data Team, Greensboro, NC 27401-4901 USA.
- Weakley, Alan S. Flora of the Southern and Mid-Atlantic States. University of North Carolina Herbarium, Working draft of 30 November 2012.
- Weakley, Alan S., Ludwig, J. Christopher and Townsend, John F. 2012. Flora of Virginia. Bland Crowder ed. Foundation of the Flora of Virginia Project, Inc., Richmond. Fort Worth: Botanical Research Institute of Texas Press.
- Wunderlin, Richard P. Guide to the Vascular Plants of Central Florida. University Press of Florida, 1982.
- Yatskievych, K. Field Guide to Indiana Wildflowers. Indiana University Press. 2000.

ERNST SEEDS WOULD LIKE TO THANK:

- The Xerces Society for Invertebrate Research and Anne Stine, Ph.D. for providing the pollinator ratings used in this catalog.
- Harland Patch, Ph.D., Assistant Research Professor, Pennsylvania State University, Dept. of Agricultural Sciences; Research Associate, Entomology at Penn State's Center for Pollinator Research.
- Dave Biddinger, Ph.D., Tree Fruit Research Entomologist and Research Professor, Pennsylvania State University, Fruit Research & Extension Center.
- Missouri Botanical Garden Plant Finder Tool.

PHOTOGRAPHS CONTRIBUTED BY THE FOLLOWING ERNST SEEDS' EMPLOYEES:

Andy Ernst	Marcia Ernst	Ayshea Heckman	Nikki Hindle
Dan Arnett	Mark Fiely	Greg Kedzierski	

Will Adams, Allison Walters Daisley, Sammie Boyer, Ben Crowl, Luke Fiely, Shayna Lantz, Sydney Winter, and Aarick Zook for assistance with data collection.

Photos contributed by outside sources are gratefully acknowledged on images or in captions throughout the catalog.

your notes

your notes

your notes

your notes

It is our honor
to serve you!

We are proud of the reputation we've built and value the relationships we've cultivated with our customers for more than half a century. We look forward to serving your ecological seed needs for many years to come.





LET'S GET
growing

Six decades of farming have taught us that working hard is the only way to work. We rise early, work late, and refuse to cut corners. Our unwavering commitment to the environment ensures we harvest, condition, and test under the highest quality standards.

Whether you are working to improve your own backyard or a sprawling corporate campus, rest easy knowing our team of pros is your team of pros. Reach out today and let's get growing.



1964 - 2024

